



## INTERVALL-HYPOXIE-TRAINING Teil 1: Theoretische Grundlagen.

Dr. med. Egor Egorov, 188 Seiten Softcover, 79,90 Euro,  
egoro Verlag, ISBN 978-3-9822267-4-3.

Mehr Informationen über den Autor und das Buch:  
[www.doc-egorov.com](http://www.doc-egorov.com)

# Literaturverzeichnis

Aboouf, M. A., Thiersch, M., Soliz, J., Gassmann, M., & Schneider Gasser, E. M. (2023). The brain at high altitude: From molecular signaling to cognitive performance. *International Journal of Molecular Sciences*, 24(12), 10179.

Aeberli, I., Erb, A., Spliethoff, K., Meier, D., Götze, O., Frühauf, H., Fox, M., Finlayson, G. S., Gassmann, M., Berneis, K., Maggiorini, M., Langhans, W., & Lutz, T. A. (2013). Disturbed eating at high altitude: Influence of food preferences, acute mountain sickness and satiation hormones. *European Journal of Nutrition*, 52(2), 625–635.

Agadzhanyan, N.A., Dvoenosov, V.G. Age-related features of the cardiorespiratory system response to the combined effects of hypoxia and hypercapnia. *General Resuscitation*. 2005;1 (2):40–44.

Ahramiyanpour, N., Shafie'ei, M., Sarvipour, N., Amiri, R., Akbari, Z. Carboxytherapy in dermatology: A systematic review. *J. Cosmet. Dermatol.* 2022, 21, 1874–1894.

Akahane, S., Sakai, Y., Ueha, T., Nishimoto, H., Inoue, M., Niikura, T., Kuroda, R. Transcutaneous carbon dioxide application accelerates muscle injury repair in rat models. *Int. Orthop.* 2017, 41, 1007–1015.

Akamine, T., Taguchi, N. Effects of an artificially carbonated bath on athletic warm-up. *J. Hum. Ergol.* 1998, 27, 22–29.

Akça, O., Doufas, A.G., Morioka, N., Iscoe, S., Fisher, J., Sessler, D.I. Hypercapnia improves tissue oxygenation. *Anesthesiology* 2002;97:801-806.

Akça, O., Sessler, D.I., Delong, D., Keijner, R., Ganzel, B., Doufas, A.G. Tissue oxygenation response to mild hypercapnia during cardiopulmonary bypass with constant pump output. *British J Anaes* 2006;96:708-714.

Alex, R. M., Panza, G. S., Hakim, H., Badr, M. S., Edwards, B. A., Sands, S. A., & Mateika, J. H. (2019). Exposure to mild intermittent hypoxia increases loop gain and the arousal threshold in participants with obstructive sleep apnoea. *The Journal of Physiology*, 597(14), 3697–3711.

Almohanna, A. M., & Wray, S. (2018). Hypoxic conditioning in blood vessels and smooth muscle tissues: Effects on function, mechanisms, and unknowns. *American Journal of Physiology. Heart and Circulatory Physiology*, 315(4), H756–H770.

Amann, M., Romer, L. M., Subudhi, A. W., Pegelow, D. F., & Dempsey, J. A. (2007). Severity of arterial hypoxaemia affects the relative contributions of peripheral muscle fatigue to exercise performance in healthy humans. *The Journal of Physiology*, 581(1), 389–403.

Amano-Iga, R., Hasegawa, T., Takeda, D., Murakami, A., Yatagai, N., Saito, I., Arimoto, S., Kakei, Y., Sakakibara, A., Akashi, M. Local application of transcutaneous carbon dioxide paste decreases inflammation and accelerates wound healing. *Cureus* 2021, 13, e19518.

Amin, N., Chen, S., Ren, Q., Tan, X., Botchway, B. O. A., Hu, Z., Chen, F., Ye, S., Du, X., Chen, Z., & Fang, M. (2021). Hypoxia inducible factor-1 $\alpha$  attenuates ischemic brain damage by modulating inflammatory response and glial activity. *Cells*, 10(6), 1359.

An, W. G., Kanekal, M., Simon, M. C., Maltepe, E., Blagosklonny, M. V., & Neckers, L. M. (1998). Stabilization of wild-type p53 by hypoxia-inducible factor 1alpha. *Nature*, 392(6674), 405–408.

Arany, Z., Foo, S., Ma, Y., Ruas, J.L., Bommi-Reddy, A., Girnun, G., Cooper, M., Laznik, D., Chinsomboon, J., Rangwala, S.M., et al. HIF-independent regulation of VEGF and angiogenesis by the transcriptional coactivator PGC-1 $\alpha$ . *Nature* 2008, 451, 1008–1012.

Arkhipenko, Y., Vdovina, I., Kostina, N., Sazontova, T., Glazachev, O. Adaptation to interval hypoxia-hyperoxia improves exercise tolerance in professional athletes: experimental substantiation and applied approbation. *European Scientific Journal*, 2014. -v.10.-P. 135-154.

Asadi, A.K., Sá, R.C., Arai, T.J., Theilmann, R.J., Hopkins, S.R., Buxton, R.B., Prisk, G.K. Regional pulmonary perfusion patterns in humans are not significantly altered by inspiratory hypercapnia. *J. Appl. Physiol.* 2019, 127, 365–375.

Baik, A. H., & Jain, I. H. (2020). Turning the oxygen dial: Balancing the highs and lows. *Trends in Cell Biology*, 30(7), 516–536.

Bailey, D. M., Bain, A. R., Hoiland, R. L., Barak, O. F., Drvis, I., Hirtz, C., Lehmann, S., Marchi, N., Janigro, D., Macleod, D. B., Ainslie, P. N., & Dujic, Z. (2022). Hypoxemia increases blood-brain barrier permeability during extreme apnea in humans. *Journal of Cerebral Blood Flow and Metabolism*, 42(6), 1120–1135.

Baird, N. A., Turnbull, D. W., & Johnson, E. A. (2006). Induction of the heat shock pathway during hypoxia requires regulation of heat shock factor by hypoxia-inducible factor-1. *The Journal of Biological Chemistry*, 281(50), 38675–38681.

Batie, M., Frost, J., Frost, M., Wilson, J. W., Schofield, P., & Rocha, S. (2019). Hypoxia induces rapid changes to histone methylation and reprograms chromatin. *Science*, 363(6432), 1222–1226.

Barer, G.R., Shaw, J.W. Pulmonary vasodilator and vasoconstrictor actions of carbon dioxide. *J. Physiol.* 1971, 213, 633–645.

Barth, A., Bauer, R., Gedrange, T., et al (1998) Influence of hypoxia and hypoxia/hypercapnia upon brain and blood peroxidative and glutathione status in normal weight and growth-restricted newborn piglets. *Exp Toxicol Pathol* 50(4-6):402–410. [https://doi.org/10.1016/S0940-2993\(98\)80026-2](https://doi.org/10.1016/S0940-2993(98)80026-2).

Baudouin, S.V., Evans, T.W. Action of carbon dioxide on hypoxic pulmonary vasoconstriction in the rat lung: Evidence against specific endothelium-derived relaxing factor-mediated vasodilation. *Crit Care Med.* 1993, 21, 740–746.

- Bayer, U., Likar, R., Pinter, G., Stettner, H., Demschar, S., Trummer, B., Neuwersch, S., Glazachev, O., & Burtscher, M. (2017). Intermittent hypoxic-hyperoxic training on cognitive performance in geriatric patients. *Alzheimer's & Dementia*, 3, 114–122.
- Bedu, M., Cheynel, J., Gascard, J., Coudert, J. Transcutaneous CO<sub>2</sub> diffusion comparison between CO<sub>2</sub> spa water and dry gas in royal thermal spa, in *Advances in Vascular Pathology*. Edited by Strano A, Novo S. Elsevier Science Publishers B. V. (Biomedical Division), 1989, pp 1109–1113.
- Behrendt, T., Bielitzki, R., Behrens, M., Herold, F., & Schega, L. (2022). Effects of intermittent hypoxia-hyperoxia on performance- and health-related outcomes in humans: A systematic review. *Sports Medicine Open*, 8(1), 70.
- Benderro, G. F., Sun, X., Kuang, Y., & Lamanna, J. C. (2012). Belaïdi, E., Ramond, A., Joyeux Faure, M. Contrasting effects of intermittent hypoxia on myocardial ischemic tolerance Intermittent hypoxia: from molecular mechanisms to clinical applications. Nova Science Publishers. 2009;3–18.
- Bellien, J., Favre, J., Iacob, M., Gao, J., Thuillez, C., Richard, V., Joannidès, R. Arterial stiffness is regulated by nitric oxide and endothelium-derived hyperpolarizing factor during changes in blood flow in humans. *Hypertension* 2010, 55, 674–680.
- Benderro, G.F., Sun, X., Kuang, Y., Lamanna, J.C. Decreased VEGF expression and microvascular density, but increased HIF-1 and 2alpha accumulation and EPO expression in chronic moderate hyperoxia in the mouse brain. *Brain Research*, 1471, 46–55.
- Benoit, H., Germain, M., Barthelemy, J.C. Preacclimatization to high altitude using exercise with normobaric hypoxic gas mixtures // *Int. J. Sports Med.* - 1992. -Vol. 13, Suppl. 1. - P. 213–216.
- Bergeron, M.f, Bahr, R., Bärtsch, P., Bourdon, L., Calbet, J., Carlsen, K.h, Castagna, O., González-Alonso, J., Lundby, C., Maughan, R.j, Millet, G., Mountjoy, M., Racinais, S., Rasmussen, P., Subudhi, A.w, Young, A.j, Soligard, T., & Engebretsen, L. (2012). International Olympic Committee consensus statement on thermoregulatory and altitude challenges for high-level athletes. *British Journal of Sports Medicine*, 46(11), 770–779.
- Bespalov, A.G., Tregub, P.P., Kulikov, V.P., et al (2014) The role of VEGF, HSP-70 and protein S-100β in the potentiation effect of the neuroprotective effect of hypercapnic hypoxia. *Patol Fiziol Eksp Ter* 2:24–27.
- Bestavashvili, A., Glazachev, O., Bestavashvili, A., Ines, D., Suvorov, A. Y., Vorontsov, N., Tuter, D., Gognieva, D., Yong, Z., & Pavlov, C. (2021). The effects of intermittent hypoxic–hyperoxic exposures on lipid profile and inflammation in patients with metabolic syndrome. *Frontiers in Cardiovascular Medicine*, 8, 700826.
- Bissonnette, S., Vaillancourt, M., Hébert, S. S., Drolet, G., & Samadi, P. (2013). Striatal pre-enkephalin overexpression improves Huntington's disease symptoms in the R6/2 mouse model of Huntington's disease. *PLoS ONE*, 8(9), e75099.
- Blackford, A. N., & Jackson, S. P. (2017). ATM, ATR, and DNA-PK: The trinity at the heart of the DNA damage response. *Molecular Cell*, 66(6), 801–817.
- Bloch, K. E., Buentzli, J. C., Latshang, T. D., & Ulrich, S. (2015). Sleep at high altitude: Guesses and facts. *Journal of Applied Physiology*, 119(12), 1466–1480.
- Bochorishvili, G., Stornetta, R. L., Coates, M. B., & Guyenet, P. G. (2012). Pre-Bötzinger complex receives glutamatergic innervation from galaninergic and other retrotrapezoid nucleus neurons. *Journal of Comparative Neurology*, 520(5), 1047–1061.
- Bohr, C., Hasselbalch, K., Krogh, A. Ueber einen in biologischer beziehung wichtigen einfluss, den die kohlensäurespannung des blutes auf dessen sauerstoffbindung übt. *Skandinavisches Archiv Für Physiologie* 1904, 16, 402–412.
- Bolevich, S., Kogan, A.; Zivkovic, V., Djuric, D., Novikov, A., Vorobyev, S., Jakovljevic, V. Protective role of carbon dioxide (CO<sub>2</sub>) in generation of reactive oxygen species. *Mol. Cell Biochem.* 2016, 411, 317–330.
- Bolland, H., Ma, T. S., Ramlee, S., Ramadan, K., & Hammond, E. M. (2021). Links between the unfolded protein response and the DNA damage response in hypoxia: A systematic review. *Biochemical Society Transactions*, 49(3), 1251–1263.
- Borukaeva, I.Kh. The pathophysiological substantiation grounds for the use of interval hypoxic training and enteral oxygenotherapy in bronchial asthma. PhD thesis. 2011;5.
- Botusan, I.R., Sunkari, V.G., Savu, O., Catrina, A.I., Grünler, J., Lindberg, S., Pereira, T., Ylä-Herttuala, S., Poellinger, L., Brismar, K., et al. Stabilization of HIF-1α is critical to improve wound healing in diabetic mice. *Proc. Nat. Acad. Sci. USA* 2008, 105, 19426–19431.
- Brown, K. A., Laferrière, A., Lakheeram, I., & Moss, I. R. (2006). Recurrent hypoxemia in children is associated with increased analgesic sensitivity to opiates. *Anesthesiology*, 105(4), 665–669.
- Burtscher, J., Bean, C., Zangrandi, L., Kmiec, I., Agostinho, A., Scorrano, L., Gnaiger, E., & Schwarzer, C. (2018). Proenkephalin derived peptides are involved in the modulation of mitochondrial respiratory control during epileptogenesis. *Frontiers in Molecular Neuroscience*, 11, 351.
- Burtscher, J., Hohenauer, E., Burtscher, M., Millet, G. P., & Egg, M. (2023). Environmental and behavioral regulation of HIF-mitochondria crosstalk. *Free Radical Biology & Medicine*, 206, 63–73.
- Burtscher, J., Maglione, V., Di Pardo, A., Millet, G. P., Schwarzer, C., & Zangrandi, L. (2021). A rationale for hypoxic and chemical conditioning in Huntington's disease. *International Journal of Molecular Sciences*, 22(2), 582.
- Burtscher, J., Mallet, R. T., Burtscher, M., & Millet, G. P. (2021). Hypoxia and brain aging: Neurodegeneration or neuroprotection? *Ageing Research Reviews*, 68, 101343.
- Burtscher, J., Mallet, R. T., Pialoux, V., Millet, G. P., & Burtscher, M. (2022). Adaptive responses to hypoxia and/or hyperoxia in humans. *Antioxidants & Redox Signaling*, 37(13–15), 887–912.
- Burtscher, J., Millet, G. P., Place, N., Kayser, B., & Zanou, N. (2021). The muscle-brain axis and neurodegenerative diseases: The key role of mitochondria in exercise-induced neuroprotection. *International Journal of Molecular Sciences*, 22(12), 6479.

- Burtscher, J., Niedermeier, M., Hüfner, K., Van Den Burg, E., Kopp, M., Stoop, R., Burtscher, M., Gatterer, H., & Millet, G. P. (2022). The interplay of hypoxic and mental stress: Implications for anxiety and depressive disorders. *Neuroscience and Biobehavioral Reviews*, 138, 104718.
- Burtscher, J., Pepe, G., Maharjan, N., Riguet, N., Di Pardo, A., Maglione, V., & Millet, G. P. (2023). Sphingolipids and impaired hypoxic stress responses in Huntington disease. *Progress in Lipid Research*, 90, 101224.
- Burtscher, J., Romani, M., Bernardo, G., Popa, T., Ziviani, E., Hummel, F. C., Sorrentino, V., & Millet, G. P. (2022). Boosting mitochondrial health to counteract neuro-degeneration. *Progress in Neurobiology*, 215, 102289.
- Burtscher, J., Syed, M. M. K., Lashuel, H. A., & Millet, G. P. (2021). Hypoxia conditioning as a promising therapeutic target in Parkinson's Disease? *Movement Disorders*, 36(4), 857–861.
- Burtscher, M., Gatterer, H., Szubski, C., Pierantozzi, E., & Faulhaber, M. (2010). Effects of interval hypoxia on exercise tolerance: Special focus on patients with CAD or COPD. *Sleep and Breathing*, 14(3), 209–220.
- Burtscher, M., Haider, T., Domej, W., Linser, T., Gatterer, H., Faulhaber, M., Pocecco, E., Ehrenburg, I., Tkatchuk, E., Koch, R., & Bernardi, L. (2009). Intermittent hypoxia increases exercise tolerance in patients at risk for or with mild COPD. *Respiratory Physiology & Neurobiology*, 165(1), 97–103.
- Burtscher, M., Mairer, K., Wille, M., Gatterer, H., Ruedl, G., Faulhaber, M. et al. Short-term exposure to hypoxia for work and leisure activities in health and disease: which level of hypoxia is safe? *Sleep Breath*. 2012;16(2):435–442.
- Burtscher, M., Pachinger, O., Ehrenbourg, I., Mitterbauer, G., Faulhaber, M., Pühringer, R., & Tkatchouk, E. (2004). Intermittent hypoxia increases exercise tolerance in elderly men with and without coronary artery disease. *International Journal of Cardiology*, 96(2), 247–254.
- Butt, U. J., Steixner-Kumar, A. A., Depp, C., Sun, T., Hassouna, I., Wüstefeld, L., Arinrad, S., Zillmann, M. R., Schopf, N., Fernandez Garcia-Agudo, L., Mohrmann, L., Bode, U., Ronnenberg, A., Hindermann, M., Goebels, S., Bonn, S., Katschinski, D. M., Miskowiak, K. W., Nave, K.-A., & Ehrenreich, H. (2021). Hippocampal neurons respond to brain activity with functional hypoxia. *Molecular Psychiatry*, 26(6), 1790–1807.
- Caldwell, H.G., Hoiland, R.L., Bain, A.R., Howe, C.A., Carr, J.M.J.R., Gibbons, T.D., Durrer, C.G., Tymko, M.M., Stacey, B.S., Bailey, D.M., Sekhon, M.S., MacLeod, D.B., Ainslie, P.N. Evidence for direct CO<sub>2</sub>-mediated alterations in cerebral oxidative metabolism in humans, July 2024, *Acta Physiologica*, DOI: 10.1111/apha.14197.
- Camacho-Cardenosa, A., Camacho-Cardenosa, M., Martínez-Guardado, I., Leal, A., Andrada, J. M. V., & Timón, R. (2022). Resistance circuit training combined with hypoxia stimulates bone system of older adults: A randomized trial. *Experimental Gerontology*, 169, 111983.
- Camacho-Cardenosa, M., Quesada-Gómez, J. M., Camacho-Cardenosa, A., Leal, A., Dorado, G., Torrecillas-Baena, B., & Casado-Díaz, A. (2020). Effects of normobaric cyclic hypoxia exposure on mesenchymal stem-cell differentiation—pilot study on bone parameters in elderly. *World Journal of Stem Cells*, 12(12), 1667–1690.
- Cantagrel, S., Krier C., Ducrocq S. et al (2003) Hypoxic preconditioning reduces apoptosis in a rat model of immature brain hypoxia-ischemia. *Neurosci Lett* 347(2):106–110.
- Carr, J. M. J. R., Caldwell, H. G., & Ainslie, P. N. (2021). Cerebral blood flow, cerebrovascular reactivity and their influence on ventilatory sensitivity. *Experimental Physiology*, 106(7), 1425–1448.
- Carr, P., Graves, J.E., Poston, L. Carbon dioxide induced vasorelaxation in rat mesenteric small arteries precontracted with noradrenaline is endothelium dependent and mediated by nitric oxide. *Pflügers Arch.* 1993, 423, 343–345.
- Cartwright, I. M., & Colgan, S. P. (2023). The hypoxic tissue microenvironment as a driver of mucosal inflammatory resolution. *Frontiers in Immunology*, 14, 1124774.
- Case, R., Greenberg, H., Moskowitz, R. Alterations in coronary sinus pO<sub>2</sub> and O<sub>2</sub> saturation resulting from pCO<sub>2</sub> changes. *Cardiovasc. Res.* 1975, 9, 167–177.
- Casey, D. P., Madery, B. D., Curry, T. B., Eisenach, J. H., Wilkins, B. W., & Joyner, M. J. (2010). Nitric oxide contributes to the augmented vasodilatation during hypoxic exercise. *The Journal of Physiology*, 588(2), 373–385.
- Chacaroun, S., Borowik, A., Morrison, S. A., Baillieul, S., Flore, P., Doutreleau, S., & Verges, S. (2016). Physiological responses to two hypoxic conditioning strategies in healthy subjects. *Frontiers in Physiology*, 7, 675.
- Chakraborty, A. A., Laukka, T., Myllykoski, M., Ringel, A. E., Booker, M. A., Tolstorukov, M. Y., Meng, Y. J., Meier, S. R., Jennings, R. B., Creech, A. L., Herbert, Z. T., Mcbrayer, S. K., Olenchock, B. A., Jaffe, J. D., Haigis, M. C., Beroukhim, R., Signoretti, S., Koivunen, P., & Kaelin, W. G. (2019). Histone demethylase KDM6A directly senses oxygen to control chromatin and cell fate. *Science*, 363(6432), 1217–1222.
- Chang, A. J., Ortega, F. E., Riegler, J., Madison, D. V., & Krasnow, M. A. (2015). Oxygen regulation of breathing through an olfactory receptor activated by lactate. *Nature*, 527(7577), 240–244.
- Charter, M.E., Lamb, I.R., Murrant, C.L. Arteriolar and capillary responses to CO<sub>2</sub> and H<sup>+</sup> in hamster skeletal muscle microvasculature: Implications for active hyperemia. *Microcirculation* 2018, 25, e12494.
- Chen, C.Y., Tsa, Y.L., Kao, C.L., Lee, S.D., Wu, M.C., Mallikarjuna K., et al. Effect of mild intermittent hypoxia on glucose tolerance, muscle morphology and AMPK\_PGC\_1alpha signaling. *Chin J Physiol.* 2010;53(1):62–71.
- Chen, F., Qi, Z., Luo, Y., Hinchliffe, T., Ding, G., Xia, Y., & Ji, X. (2014). Non-pharmaceutical therapies for stroke: Mechanisms and clinical implications. *Progress in Neurobiology*, 115, 246–269.
- Chen, T., Li, J., Chao, D., Sandhu, H. K., Liao, X., Zhao, J., Wen, G., & Xia, Y. (2014). δ-Opioid receptor activation reduces α-synuclein overexpression and oligomer formation induced by MPP<sup>+</sup> and/or hypoxia. *Experimental Neurology*, 255, 127–136.

- Chen, W.J., Chen, H.W., Yu, S.L., et al (2005) Gene expression profiles in hypoxic preconditioning using cDNA microarray analysis: altered expression of an angiogenic factor, carcinoembryonic antigen-related cell adhesion molecule 1. Shock 24:124–131.
- Chen, Y., Cann, M.J., Litvin, T.N., et al. Soluble Adenylyl Cyclase as an Evolutionarily Conserved Bicarbonate Sensor. Science 2000;289:625–628.
- Cheng, S.-C., Quintin, J., Cramer, R. A., Shepardson, K. M., Saeed, S., Kumar, V., Giamparellos-Bourboulis, E. J., Martens, J. H. A., Rao, N. A., Aghajanirefah, A., Manjeri, G. R., Li, Y., Ifrim, D. C., Arts, R. J. W., Van Der Veer, B. M. J. W., Deen, P. M. T., Logie, C., O’Neill, L. A., Willems, P., Van De Veerdonk, F. L., Van Der Meer, J. W. M., Ng, A., Joosten, L. A. B., Wijmenga, C., Stunnenberg, H. G., Xavier, R. J., & Netea, M. G. (2014). mTOR-and HIF-1 $\alpha$ -mediated aerobic glycolysis as metabolic basis for trained immunity. Science, 345(6204), 1250684.
- Chudimov, V.F., Kulikov, V.P., Kuropatyatnik, N.I., Boyko, E.A., Shadrina, E.N., Voronkova, E.I. The use of hypoxic-hypercapnic training in children with attention deficit hyperactivity disorder for correcting the problems of school disadaptation. Voprosy balneologii, fizioterapii i lechebnoi fizicheskoi kultury. = Questions of Balneology, Physiotherapy and Therapeutic Physical Training. 2011;147(3):36–39.
- Chudimov, V.F., Kulikov, V.P., Kuropatyatnik, N.I., Boyko, E.A., Voronkova, E.I. The influence of hypoxic-hypercapnic training on the well-being, activity, mood and performance of schoolchildren, classified for health reasons to a special medical group for physical education. Lechebnaia Fizkultura i Sportivnaya Meditsina = Therapeutic Physical Training and Sports Medicine. 2010;80(8):45–50.
- Clanton, T. L., & Klawitter, P. F. (2001). Invited review: Adaptive responses of skeletal muscle to intermittent hypoxia: the known and the unknown. Journal of Applied Physiology, 90(6), 2476–2487.
- Contreras, M., Masterson, C., Laffey, J.G. Permissive hypercapnia: What to remember. Curr. Opin. Anaesthesiol. 2015, 28, 26–37.
- Coskun, B., Aras, D., Akalan, C., Kocak, S., & Hamlin, M. J. (2022). Plyometric Training in normobaric hypoxia improves jump performance. International Journal of Sports Medicine, 43, 519–525.
- Cummins, E.P., Oliver, K.M., Lenihan, C.R., Fitzpatrick, S.F., Brunning, U., Scholz, C.C., Slattery, C., Leonard, M.O., McLoughlin, P., Taylor, C.T. NF- $\kappa$ B links CO<sub>2</sub> sensing to innate immunity and inflammation in mammalian cells. J. Immunol. 2010, 185, 4439–4445.
- Crystal, G.J. Carbon dioxide and the heart: Physiology and clinical implications. Anesth. Analg. 2015, 121, 610–623.
- D’Arcangelo, D., Facchiano, F., Barlucchi, L., et al. Acidosis inhibits endothelial cell apoptosis and function and induces basic fibroblast growth factor and vascular endothelial growth factor expression. Circ Res 2000;86:312–318.
- D’hulst, G., & Deldicque, L. (2017). Human skeletal muscle wasting in hypoxia: A matter of hypoxic dose? Journal of Applied Physiology, 122(2), 406–408.
- D’ignazio, L., Bandarra, D., & Rocha, S. (2016). NF- $\kappa$ B and HIF crosstalk in immune responses. FEBS Journal, 283(3), 413–424.
- D’Ignazio, L., & Rocha, S. (2016). Hypoxia Induced NF- $\kappa$ B. Cells, 5, 10.
- De Kloet, E. R. (2013). Functional profile of the binary brain corticosteroid receptor system: Mediating, multitasking, coordinating, integrating. European Journal of Pharmacology, 719(1–3), 53–62.
- Deacon, N. L., McEvoy, R. D., Stadler, D. L., & Catcheside, P. G. (2017). Intermittent hypercapnic hypoxia during sleep does not induce ventilatory long-term facilitation in healthy males. Journal of Applied Physiology, 123(3), 534–543.
- Deng, M., Yang, X., Qin, B., Liu, T., Zhang, H., Guo, W., Lee, S., Kim, J., Yuan, J., Pei, H., et al. Deubiquitination and activation of AMPK by USP10. Mol. Cell. 2016, 61, 614–624.
- Dengler, F. (2020). Activation of AMPK under hypoxia: Many roads leading to Rome. International Journal of Molecular Sciences, 21(7), 2428.
- Diebold, I., Flugel, D., Becht, S., Belaiba, R. S., Bonello, S., Hess, J., Kietzmann, T., & Gorlach, A. (2010). The hypoxia-inducible factor-2alpha is stabilized by oxidative stress involving NOX4. Antioxidants & Redox Signaling, 13, 425–436.
- Diebold, I., Petry, A., Hess, J., & Görlich, A. (2010). The NADPH oxidase subunit NOX4 is a new target gene of the hypoxia-inducible factor-1. Molecular Biology of the Cell, 21(12), 2087–2096.
- Ding, F.-S., Cheng, X., Zhao, T., Zhao, Y.-Q., Zhang, G.-B., Wu, H.-T., Zhu, L.-L., & Wu, K.-W. (2019). [Intermittent hypoxic preconditioning relieves fear and anxiety behavior in post-traumatic stress model mice]. Sheng Li Xue Bao [Acta Physiologica Sinica], 71, 537–546.
- Dirckx, N., Tower, R. J., Mercken, E. M., Vangoitsenhoven, R., Moreau-Triby, C., Breugelmans, T., Nefyodova, E., Cardoen, R., Mathieu, C., Van Der Schueren, B., Confavreux, C. B., Clemens, T. L., & Maes, C. (2018). Vhl deletion in osteoblasts boosts cellular glycolysis and improves global glucose metabolism. Journal of Clinical Investigation, 128(3), 1087–1105.
- Dirnagl, U., Becker, K., & Meisel, A. (2009). Preconditioning and tolerance against cerebral ischaemia: From experimental strategies to clinical use. The Lancet Neurology, 8(4), 398–412.
- Dogliotti, G., Galliera, E., Iorio, E., De Bernardi Di Valserra, M., Solimene, U., Corsi, M.M. Effect of immersion in CO<sub>2</sub>-enriched water on free radical release and total antioxidant status in peripheral arterial occlusive disease. Int. Angiol. 2011, 30, 12–17.
- Dorrington, K.L., Talbot, N.P. Human pulmonary vascular responses to hypoxia and hypercapnia. Pflugers Arch. 2004, 449, 1–15.
- Dospinescu, V., Nijjar, S., Spanos, F., Cook, J., de Wolf, E., Biscotti, M.A., Gerdol, M., Dale, N. Structural determinants of CO<sub>2</sub>-sensitivity in the  $\beta$  connexin family suggested by evolutionary analysis. Commun. Biol. 2019, 2, 331.

- Dudnik, E., Zagaynaya, E., Glazachev, O. S., & Susta, D. (2018). Intermittent hypoxia-hyperoxia conditioning improves cardiorespiratory fitness in older comorbid cardiac outpatients without hematological changes: A randomized controlled trial. *High Altitude Medicine & Biology*, 19, 339–343.
- Duling, B.R. Changes in microvascular diameter and oxygen tension induced by carbon dioxide. *Circ. Res.* 1973, 32, 370–376.
- Ehrenreich, H., Garcia-Agudo, L. F., Steixner-Kumar, A. A., Wilke, J. B. H., & Butt, U. J. (2022). Introducing the brain erythropoietin circle to explain adaptive brain hardware upgrade and improved performance. *Molecular Psychiatry*, 27(5), 2372–2379.
- Ehrenreich, H., Gassmann, M., Poustka, L., Burtscher, M., Hammermann, P., Sirén, A.-L., Nave, K.-A., & Miskowiak, K. (2023). Exploiting moderate hypoxia to benefit patients with brain disease: Molecular mechanisms and translational research in progress. *Neuroprotection*, 1(1), 9–19.
- El-Chami, M., Sudan, S., Lin, H.-o.-S., & Mateika, J. H. (2017). Exposure to intermittent hypoxia and sustained hypercapnia reduces therapeutic CPAP in participants with obstructive sleep apnea. *Journal of Applied Physiology*, 123(4), 993–1002.
- Eltzschig, H. K., & Carmeliet, P. (2011). Hypoxia and inflammation. *New England Journal of Medicine*, 364(7), 656–665.
- Esparza-Moltó, P. B., Romero-Carramiñana, I., Núñez De Arenas, C., Pereira, M. P., Blanco, N., Pardo, B., Bates, G. R., Sánchez-Castillo, C., Artuch, R., Murphy, M. P., Esteban, J. A., & Cuevza, J. M. (2021). Generation of mitochondrial reactive oxygen species is controlled by ATPase inhibitory factor 1 and regulates cognition. *PLoS Biology*, 19(5), e3001252.
- Estrada, J. A., Williams, A. G., Sun, J., Gonzalez, L., Downey, H. F., Caffrey, J. L., & Mallet, R. T. (2016). delta-Opioid receptor (DOR) signaling and reactive oxygen species (ROS) mediate intermittent hypoxia induced protection of canine myocardium. *Basic Research in Cardiology*, 111(2), 17.
- Etulain, J., Negrotto, S., Carestia, A., Pozner, R., Romaniuk, M., D'atri, L., Klement, G., & Schattner, M. (2012). Acidosis downregulates platelet haemostatic functions and promotes neutrophil proinflammatory responses mediated by platelets. *Thrombosis and Haemostasis*, 107(01), 99–110.
- Fabry, R., Monnet, P., Schmidt, J., Lusson, J.R., Carpentier, P-H., Baguet, J-C,Dubray, C. Clinical and microcirculatory effects of transcutaneous CO<sub>2</sub> therapy in intermittent claudication. Randomized double-blind clinical trial with a parallel design. *Vasa* 2009, 38, 213–224.
- Fernández-Agüera, M. C., Gao, L., González-Rodríguez, P., Pintado, C. O., Arias-Mayenco, I., García-Flores, P., García-Pergañeda, A., Pascual, A., Ortega-Sáenz, P., & López-Barneo, J. (2015). Oxygen sensing by arterial chemoreceptors depends on mitochondrial complex I signaling. *Cell Metabolism*, 22(5), 825–837.
- Fernández-Lázaro, D., Mielgo-Ayuso, J., Adams, D. P., González-Bernal, J. J., Fernández Araque, A., Cano García, A., & Fernández-Lázaro, C. I. (2020). Electromyography: A simple and accessible tool to assess physical performance and health during hypoxia training. A systematic review. *Sustainability*, 12(21), 9137.
- Ferrer, I. (2006) Apoptosis: future targets for neuroprotective strategies. *Cerebrovasc Dis* 21(Suppl 2):9–20. <https://doi.org/10.1159/000091699>.
- Fesenko, M.S., Lisyana, T.O. Hypoxic stimulation in prolonged and recurrent bronchitis in young children. *Physiology Journal*. 1992;38 (5):31–33.
- Forster, H. V., & Smith, C. A. (2010). Contributions of central and peripheral chemoreceptors to the ventilatory response to CO<sub>2</sub>/H<sup>+</sup>. *Journal of Applied Physiology*, 108(4), 989–994.
- Foskolou, I. P., Jorgensen, C., Leszczynska, K. B., Olcina, M. M., Tarhonskaya, H., Haisma, B., D'angioloella, V., Myers, W. K., Domene, C., Flashman, E., & Hammond, E. M. (2017). Ribonucleotide reductase requires subunit switching in hypoxia to maintain DNA replication. *Molecular Cell*, 66(2), 206–220.e9.e209.
- Foster, G. E., Brugniaux, J. V., Pialoux, V., Duggan, C. T. C., Hanly, P. J., Ahmed, S. B., & Poulin, M. J. (2009). Cardiovascular and cerebrovascular responses to acute hypoxia following exposure to intermittent hypoxia in healthy humans. *The Journal of Physiology*, 587(13), 3287–3299.
- Friedl, P. (2004). Prespecification and plasticity: Shifting mechanisms of cell migration. *Current Opinion in Cell Biology*, 16(1), 14–23.
- Fukuda, R., Zhang, H., Kim, J.-W., Shimoda, L., Dang, C. V., & Semenza, G. L. (2007). HIF-1 regulates cytochrome oxidase subunits to optimize efficiency of respiration in hypoxic cells. *Cell*, 129(1), 111–122.
- Funk, G. D., & Gourine, A. V. (2018). CrossTalk proposal: A central hypoxia sensor contributes to the excitatory hypoxic ventilatory response. *The Journal of Physiology*, 596(15), 2935–2938.
- Furian, M., Bitos, K., Hartmann, S. E., Muralt, L., Lichtblau, M., Bader, P. R., Rawling, J. M., Ulrich, S., Poulin, M. J., & Bloch, K. E. (2022). Acute high altitude exposure, acclimatization and re-exposure on nocturnal breathing. *Frontiers in Physiology*, 13, 965021.
- Galganska, H., Jarmuszkiewicz, W., Galganski, L. Carbon dioxide inhibits COVID-19-type proinflammatory responses through extracellular signal-regulated kinases 1 and 2, novel carbon dioxide sensors. *Cell. Mol. Life Sci.* 2021, 78, 8229–8242.
- Gangwar, A., Pooja, Sharma, M., Singh, K., Patyal, A., Bhaumik, G., Bhargava, K., & Sethy, N. K. (2019). Intermittent normobaric hypoxia facilitates high altitude acclimatization by curtailing hypoxia-induced inflammation and dyslipidemia. *Pflügers Archiv: European Journal of Physiology*, 471(7), 949–959.
- Gao, C.-J., Niu, L., Ren, P.-C., Wang, W., Zhu, C., Li, Y.-Q., Chai, W., & Sun, X.-D. (2012). Hypoxic preconditioning attenuates global cerebral ischemic injury following asphyxial cardiac arrest through regulation of delta opioid receptor system. *Neuroscience*, 202, 352–362.
- Geers, C., Gros, G. Carbon Dioxide Transport and Carbonic Anhydrase in Blood and Muscle. *Physiol Rev* 2000;80:681-715.

- Gerst, D. G., Yokhana, S. S., Carney, L. M., Lee, D. S., Badr, M. S., Qureshi, T., Anthouard, M. N., & Mateika, J. H. (2011). The hypoxic ventilatory response and ventilatory long-term facilitation are altered by time of day and repeated daily exposure to intermittent hypoxia. *Journal of Applied Physiology*, 110(1), 15–28.
- Gibson, O. R., Taylor, L., Watt, P. W., & Maxwell, N. S. (2017). Cross-adaptation: Heat and cold adaptation to improve physiological and cellular responses to hypoxia. *Sports Medicine*, 47(9), 1751–1768.
- Glazachev, O., Kopylov, P., Susta, D., Dudnik, E., & Zagaynaya, E. (2017). Adaptations following an intermittent hypoxia-hyperoxia training in coronary artery disease patients: A controlled study. *Clinical Cardiology*, 40(6), 370–376.
- Glazachev, O. S., Kryzhanovskaya, S. Y.u, Zapara, M. A., Dudnik, E. N., Samartseva, V. G., & Susta, D. (2021). Safety and efficacy of intermittent hypoxia conditioning as a new rehabilitation/secondary prevention strategy for patients with cardiovascular diseases: A systematic review and meta-analysis. *Current Cardiology Reviews*, 17(6), e051121193317.
- Godoy, L. D., Rossignoli, M. T., Delfino-Pereira, P., Garcia-Cairasco, N., & De Lima Umeoka, E. H. (2018). A comprehensive overview on stress neurobiology: Basic concepts and clinical implications. *Frontiers in Behavioral Neuroscience*, 12, 127.
- Gonchar, O., & Mankovska, I. (2012). Moderate hypoxia/hyperoxia attenuates acute hypoxia-induced oxidative damage and improves antioxidant defense in lung mitochondria. *Acta Physiologica Hungarica*, 99(4), 436–446.
- Goranchuk, V.V., Sapova, N.I., Ivanov, A.O. Hypoxic therapy. St Petersburg: Elbi-SPb, 2003. 536 p.
- Gordan, J. D., Bertout, J. A., Hu, C.-J., Diehl, J. A., & Simon, M. C. (2007). HIF-2 $\alpha$  promotes hypoxic cell proliferation by enhancing c-Myc transcriptional activity. *Cancer Cell*, 11(4), 335–347.
- Goridis, C., & Brunet, J.-F. (2010). Central chemoreception: Lessons from mouse and human genetics. *Respiratory Physiology & Neurobiology*, 173(3), 312–321.
- Goss S.P., Singh R.J., Kalyanaraman B. (1999) Bicarbonate enhances the peroxidase activity of Cu, Zn-superoxide dismutase. Role of carbonate anion radical. *J Biol Chem* 274(40):28233–28239. <https://doi.org/10.1074/jbc.274.40.28233>.
- Gourine, A. V., & Funk, G. D. (2017). On the existence of a central respiratory oxygen sensor. *Journal of Applied Physiology*, 123(5), 1344–1349.
- Guan, Y., Gu, Y., Shao, H., Ma, W., Li, G., Guo, M., Shao, Q., Li, Y., Liu, Y., Wang, C., Tian, Z., Liu, J., & Ji, X. (2023). Intermittent hypoxia protects against hypoxic-ischemic brain damage by inducing functional angiogenesis. *Journal of Cerebral Blood Flow and Metabolism*, 43(10), 1656–1671.
- Guan, Y., Liu, J., Gu, Y., & Ji, X. (2023). Effects of hypoxia on cerebral microvascular angiogenesis: Benefits or damages? *Aging and Disease*, 14(2), 370–385.
- Guner, I., Uzun, D. D., Yaman, M. O., Genc, H., Gelisgen, R., Korkmaz, G. G., Hallac, M., Yelmen, N., Sahin, G., Karter, Y., &
- Simsek, G. (2013). The effect of chronic long-term intermittent hypobaric hypoxia on bone mineral density in rats: Role of nitric oxide. *Biological Trace Element Research*, 154(2), 262–267.
- Gusarova, G. A., Dada, L. A., Kelly, A. M., Brodie, C., Witters, L. A., Chandel, N. S., & Sznajder, J. I. (2009). Alpha1-AMP-activated protein kinase regulates hypoxia-induced Na,K-ATPase endocytosis via direct phosphorylation of protein kinase C zeta. *Molecular and Cellular Biology*, 29(13), 3455–3464.
- Guyenet, P. G., Stornetta, R. L., Souza, G. M. P. R., Abbott, S. B. G., Shi, Y., & Bayliss, D. A. (2019). The retrotrapezoid nucleus: Central chemoreceptor and regulator of breathing automaticity. *Trends in Neuroscience (Tins)*, 42(11), 807–824.
- Haase, V. H. (2013). Regulation of erythropoiesis by hypoxia-inducible factors. *Blood Reviews*, 27(1), 41–53.
- Haider, T., Casucci, G., Linser, T., Faulhaber, M., Gatterer, H., Ott, G., Linser, A., Ehrenbourg, I., Tkatchouk, E., Burtscher, M., & Bernardi, L. (2009). Interval hypoxic training improves autonomic cardiovascular and respiratory control in patients with mild chronic obstructive pulmonary disease. *Journal of Hypertension*, 27(8), 1648–1654.
- Ham, P. B., & Raju, R. (2017). Mitochondrial function in hypoxic ischemic injury and influence of aging. *Progress in Neurobiology*, 157, 92–116.
- Hanly, E.J., Bachman, S.L., Marohn, M.R., Boden, J.H., Herring, A.E., De Maio, A., Talamini, M.A. Carbon dioxide pneumoperitoneum-mediated attenuation of the inflammatory response is independent of systemic acidosis. *Surgery* 2005, 137, 559–566.
- Hanly, E.J., Fuentes, J.M., Aurora, A.R., Bachman, S.L., De Maio, A., Marohn, M.R., Talamini, M.A. Carbon dioxide pneumoperitoneum prevents mortality from sepsis. *Surg. Endosc.* 2006, 20, 1482–1487.
- Hao, T., Yu, J., Wu, Z., Jiang, J., Gong, L., Wang, B., Guo, H., Zhao, H., Lu, B., Engelender, S., He, H.e, & Song, Z. (2023). Hypoxia-reprogrammed megamitochondrion contacts and engulfs lysosome to mediate mitochondrial self-digestion. *Nature Communications*, 14(1), 4105.
- Harada, R., Kawamoto, T., Ueha, T., Minoda, M., Toda, M., Onishi, Y., Fukase, N., Hara, H., Sakai, Y., Miwa, M., et al. Reoxygenation using a novel CO<sub>2</sub> therapy decreases the metastatic potential of osteosarcoma cells. *Exp. Cell Res.* 2013, 319, 1988–1997.
- Hardie, D.G., Ross, F.A., Hawley, S.A. AMPK: A nutrient and energy sensor that maintains energy homeostasis. *Nat. Rev. Mol. Cell Biol.* 2012, 13, 251–262.
- Hardie, D.G., Ross, F.A., Hawley, S.A. AMPK: A nutrient and energy sensor that maintains energy homeostasis. *Nat. Rev. Mol. Cell Biol.* 2012, 13, 251–262.
- Harris, D. P., Balasubramaniam, A., Badr, M. S., & Mateika, J. H. (2006). Long-term facilitation of ventilation and genioglossus muscle activity is evident in the presence of elevated levels of carbon dioxide in awake humans. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 291(4), R1111–R1119.

- Hartmann, B.R., Bassenge, E., Hartmann, M. Effects of serial percutaneous application of carbon dioxide in intermittent claudication: Results of a controlled trial. *Angiology* 1997, 48, 957–963.
- Haufe, S., Wiesner, S., Engeli, S., Luft, F. C., & Jordan, J. (2008). Influences of normobaric hypoxia training on metabolic risk markers in human subjects. *Medicine and Science in Sports and Exercise*, 40(11), 1939–1944.
- Hayashi, T., Asano, Y., Shintani, Y., Aoyama, H., Kioka, H., Tsukamoto, O., Hikita, M., Shinzawa-Itoh, K., Takafuji, K., Higo, S., Kato, H., Yamazaki, S., Matsuoaka, K., Nakano, A., Asanuma, H., Asakura, M., Minamino, T., Goto, Y.u-I., Ogura, T., Kitakaze, M., Komuro, I., Sakata, Y., Tsukihara, T., Yoshikawa, S., & Takashima, S. (2015). HIF1a is a positive regulator of cytochrome c oxidase. *Proceedings of the National Academy of Sciences*, 112(5), 1553–1558.
- Hayes, H. B., Jayaraman, A., Herrmann, M., Mitchell, G. S., Rymer, W. Z., & Trumbower, R. D. (2014). Daily intermittent hypoxia enhances walking after chronic spinal cord injury: a randomized trial. *Neurology*, 82(2), 104–113.
- Hill, E., Dale, N., Wall, M.J. CO<sub>2</sub>-sensitive connexin hemichannels in neurons and glia: Three different modes of signalling? *Int. J. Mol. Sci.* 2021, 22, 7254.
- Hoiland, R. L., Bain, A. R., Rieger, M. G., Bailey, D. M., & Ainslie, P. N. (2016). Hypoxemia, oxygen content, and the regulation of cerebral blood flow. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 310(5), R398–R413.
- Holmquist-Mengelbier, L., Fredlund, E., Löfstedt, T., Noguera, R., Navarro, S., Nilsson, H., Pietras, A., Vallon-Christersson, J., Borg, Å., Gradin, K., Poellinger, L., & Pahlman, S. (2006). Recruitment of HIF-1a and HIF-2a to common target genes is differentially regulated in neuroblastoma: HIF-2a promotes an aggressive phenotype. *Cancer Cell*, 10(5), 413–423.
- Hosford, P.S., Wells, J.A., Nizari, S., Christie, I.N., Theparambil, S.M., Castro, P.A., Hadjihambi, A., Barros, L.F., Ruminot, I., Lythgoe, M.F., et al. CO<sub>2</sub> signaling mediates neurovascular coupling in the cerebral cortex. *Nat. Commun.* 2022, 13, 2125.
- Hsiao, H.-T., Lin, Y.-C., Wang, J. C.-F., Tsai, Y.-C., & Liu, Y.-C. (2016). Hypoxia inducible factor-1α inhibition produced anti-allodynia effect and suppressed inflammatory cytokine production in early stage of mouse complex regional pain syndrome model. *Clinical and Experimental Pharmacology & Physiology*, 43(3), 355–359.
- Huckstepp, R.T.R., Id Bihi, R., Eason, R., et al. Connexin hemichannel-mediated CO<sub>2</sub>-dependent release of ATP in the medulla oblongata contributes to central respiratory chemosensitivity. *J Physiol* 2010;588:3901–3920.
- Inoue, S., Moriyama, H., Yakuwa, T., et al. Transcutaneous carbon dioxide improves contractures after spinal cord injury in rats. *Clin Orthop Relat Res* 2019;477:1934–1946.
- Inoue, M., Sakai, Y., Oe, K., Ueha, T., Koga, T., Nishimoto, H., Akahane, S., Harada, R., Lee, S.Y., Niikura, T., et al. Transcutaneous carbon dioxide application inhibits muscle atrophy after fracture in rats. *J. Orthop. Sci.* 2020, 25, 338–343.
- Irie, H., Tatsumi, T., Takamiya, M., Zen, K., Takahashi, T., Azuma, A., Tateishi, K., Nomura, T., Hayashi, H., Nakajima, N., et al. Carbon dioxide-rich water bathing enhances collateral blood flow in ischemic hindlimb via mobilization of endothelial progenitor cells and activation of NO-cGMP system. *Circulation* 2005, 111, 1523–1529.
- Iturriaga, R. (2023). Carotid body contribution to the physio-pathological consequences of intermittent hypoxia: Role of nitro-oxidative stress and inflammation. *The Journal of Physiology*. Advance Online Publication. <https://doi.org/10.1113/JP284112>.
- Iturriaga, R., Alcayaga, J., Chapleau, M. W., & Somers, V. K. (2021). Carotid body chemoreceptors: Physiology, pathology, and implications for health and disease. *Physiological Reviews*, 101(3), 1177–1235.
- Iwata, E., Hasegawa, T., Ueha, T., Takeda, D., Saito, I., Kawamoto, T., Akisue, T., Sakai, Y., Sasaki, R., Kuroda, R., et al. Transcutaneous carbon dioxide enhances the antitumor effect of radiotherapy on oral squamous cell carcinoma. *Oncol. Rep.* 2018, 40, 434–442.
- Iyer, N. V., Kotch, L. E., Agani, F., Leung, S. W., Laughner, E., Wenger, R. H., Gassmann, M., Gearhart, J. D., Lawler, A. M., Yu, A. Y., & Semenza, G. L. (1998). Cellular and developmental control of O<sub>2</sub> homeostasis by hypoxia-inducible factor 1 alpha. *Genes & Development*, 12(2), 149–162.
- Izumi, Y., Yamaguchi, T., Yamazaki, T., Yamashita, N., Nakamura, Y., Shiota, M., Tanaka, M., Sano, S., Osada-Oka, M., Shimada, K., et al. Percutaneous carbon dioxide treatment using a gas mist generator enhances the collateral blood flow in the ischemic hindlimb. *J. Atheroscl. Thromb.* 2015, 22, 38–51.
- Jankov, R.P., Kavanagh, B.P., Teixeira, L., et al. Therapeutic hypercapnia prevents chronic hypoxia-induced pulmonary hypertension in the newborn rat. *Amer J Physiol*. 2006;291:L912.
- Jarrard, C. P., Nagel, M. J., Stray-Gundersen, S., Tanaka, H., & Lalande, S. (2021). Hypoxic preconditioning attenuates ischemia-reperfusion injury in young healthy adults. *Journal of Applied Physiology*, 130(3), 846–852.
- Jaskiewicz, M., Moszynska, A., Króliczewski, J., Cabaj, A., Bartoszewska, S., Charzynska, A., Gebert, M., Dabrowski, M., Collawn, J. F., & Bartoszowski, R. (2022). The transition from HIF-1 to HIF-2 during prolonged hypoxia results from reactivation of PHDs and HIF1A mRNA instability. *Cellular & Molecular Biology Letters*, 27(1), 109.
- Jensen, F.B. Red blood cell pH, the Bohr effect, and other oxygenation-linked phenomena in blood O<sub>2</sub> and CO<sub>2</sub> transport. *Acta Physiologica* 2004;182:215–227.
- Jiang, B. H., Semenza, G. L., Bauer, C., & Marti, H. H. (1996). Hypoxia-inducible factor 1 levels vary exponentially over a physiologically relevant range of O<sub>2</sub> tension. *American Journal of Physiology*, 271(4), C1172–C1180.
- Jiménez-Gómez, B., Ortega-Sáenz, P., Gao, L., González-Rodríguez, P., García-Flores, P., Chandel, N., & López-Barneo, J. (2023). Transgenic NADH dehydrogenase restores oxygen regulation of breathing in mitochondrial complex I-deficient mice. *Nature Communications*, 14(1), 1172.

- Ju, X., Mallet, R. T., Downey, H. F., Metzger, D. B., & Jung, M. E. (2012). Intermittent hypoxia conditioning protects mitochondrial cytochrome c oxidase of rat cerebellum from ethanol withdrawal stress. *Journal of Applied Physiology*, 112(10), 1706–1714.
- Jugdutt, B. I. (2002). Nitric oxide and cardioprotection during ischemia-reperfusion. *Heart Failure Reviews*, 7(4), 391–405.
- Jung, M. E., Simpkins, J. W., Wilson, A. M., Downey, H. F., & Mallet, R. T. (2008). Intermittent hypoxia conditioning prevents behavioral deficit and brain oxidative stress in ethanol-withdrawn rats. *Journal of Applied Physiology*, 105(2), 510–517.
- Kaldenhoff, R., Kai, L., Uehlein, N. Aquaporins and membrane diffusion of CO<sub>2</sub> in living organisms. *Biochimica et biophysica acta* 2014;1840:1592-1595.
- Kallenborn-Gerhardt, W., Schroder, K., Del Turco, D., Lu, R., Kynast, K., Kosowski, J., Niederberger, E., Shah, A. M., Brandes, R. P., Geisslinger, G., & Schmidtko, A. (2012). NADPH oxidase-4 maintains neuropathic pain after peripheral nerve injury. *Journal of Neuroscience*, 32(30), 10136–10145.
- Kanngiesser, M., Mair, N., Lim, H. Y., Zschiesbsch, K., Blees, J., Häussler, A., Brüne, B., Ferreiròs, N., Kress, M., & Tegeder, I. (2014). Hypoxia-inducible factor 1 regulates heat and cold pain sensitivity and persistence. *Antioxidants & Redox Signaling*, 20, 2555–2571.
- Karash, Yu.M. Normobaric hypoxia in the treatment, prevention and rehabilitation. Edited by R.B. Strelkov, A.Yu. Chizhov. M.: Medicine, 1988. 352 p.
- Karpova, E.S., Kotelnikova, E.V., Lyamina, N.P. Ischemic preconditioning and its cardioprotective effect in cardiac rehabilitation programs for patients with coronary heart disease after percutaneous coronary interventions. *Ros Cardiol J.* 2012;4:104–108.
- Katayama, K., Yoshitake, Y., Watanabe, K., Akima, H., & Ishida, K. (2010). Muscle deoxygenation during sustained and intermittent isometric exercise in hypoxia. *Medicine and Science in Sports and Exercise*, 42(7), 1269–1278.
- Katayama, N., Sugimoto, K., Okada, T., Ueha, T., Sakai, Y., Akiyoshi, H., Mie, K., Ueshima, E., Sofue, K., Koide, Y., et al. Intra-arterially infused carbon dioxide-saturated solution for sensitizing the anticancer effect of cisplatin in a rabbit VX2 liver tumor model. *Int. J. Oncol.* 2017, 51, 695–701.
- Kayser, B., & Verges, S. (2013). Hypoxia, energy balance and obesity: From pathophysiological mechanisms to new treatment strategies. *Obesity Reviews*, 14(7), 579–592.
- Keir, D. A., Duffin, J., Millar, P. J., & Floras, J. S. (2019). Simultaneous assessment of central and peripheral chemoreflex regulation of muscle sympathetic nerve activity and ventilation in healthy young men. *The Journal of Physiology*, 597(13), 3281–3296.
- Keith, B., Johnson, R. S., & Simon, M. C. (2011). HIF1α and HIF2α: Sibling rivalry in hypoxic tumour growth and progression. *Nature Reviews Cancer*, 12(1), 9–22.
- Kelly, M. N., Smith, D. N., Sunshine, M. D., Ross, A., Zhang, X., Gumz, M. L., Esser, K. A., & Mitchell, G. S. (2020). Circadian clock genes and respiratory neuroplasticity genes oscillate in the phrenic motor system. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 318(6), R1058–R1067.
- Kemp, P. J., Lewis, A., Hartness, M. E., Searle, G. J., Miller, P., O'Kelly, I., & Peers, C. (2002). Airway chemotransduction: from oxygen sensor to cellular effector. *American Journal of Respiratory and Critical Care Medicine*, 166(1), S17–S24.
- Keogh, C.E., Scholz, C.C., Rodriguez, J., Selfridge, A.C., von Kriegsheim, A., Cummins, E.P. Carbon dioxide-dependent regulation of NF-kappa-B family members RelB and p100 gives molecular insight into CO<sub>2</sub>-dependent immune regulation. *J. Biol. Chem.* 2017, 292, 11561–11571.
- Khodadadeh, B., Badr, M. S., & Mateika, J. H. (2006). The ventilatory response to carbon dioxide and sustained hypoxia is enhanced after episodic hypoxia in OSA patients. *Respiratory Physiology & Neurobiology*, 150(2–3), 122–134.
- Kierans, S.J., Taylor, C.T. Regulation of glycolysis by the hypoxia-inducible factor (HIF): Implications for cellular physiology. *J. Physiol.* 2021, 599, 23–37.
- Kim, J.-W., Tchernyshyov, I., Semenza, G. L., & Dang, C. V. (2006). HIF-1-mediated expression of pyruvate dehydrogenase kinase: A metabolic switch required for cellular adaptation to hypoxia. *Cell Metabolism*, 3(3), 177–185.
- Kirby, B.S., Crecelius, A.R., Richards, J.C., Dineno, F.A. Sources of intravascular ATP during exercise in humans: Critical role for skeletal muscle perfusion. *Exp. Physiol.* 2013, 98, 988–998.
- Kivelä, R., Silvennoinen, M., Touvrä, A., Lehti, T.M., Kainulainen, H., Viiko, V. Effects of experimental type 1 diabetes and exercise training on angiogenic gene expression and capillarization in skeletal muscle. *FASEB J.* 2006, 20, 1570–1572.
- Koga, T., Niikura, T., Lee, S.Y., Okumachi, E., Ueha, T., Iwakura, T., Sakai, Y., Miwa, M., Kuroda, R., Kurosaka, M. Topical cutaneous CO<sub>2</sub> application by means of a novel hydrogel accelerates fracture repair in rats. *J. Bone Jt. Surg.* 2014, 96, 2077–2084.
- Kong, Z., Zang, Y., & Hu, Y. (2014). Normobaric hypoxia training causes more weight loss than normoxia training after a 4-week residential camp for obese young adults. *Sleep & breathing = Schlaf & Atmung*, 18(3), 591–597.
- Krook, A., Wallberg-Henriksson, H., & Zierath, J. R. (2004). Sending the signal: Molecular mechanisms regulating glucose uptake. *Medicine and Science in Sports and Exercise*, 36(7), 1212–1217.
- Kulikov, V.P., Bespalov, A.G. The device and method of creating hypercapnia for evaluation of perfusion reserve of cerebral circulation. *Echography*. 2002;3(2):160–164.
- Kulikov, V.P., Bespalov, A.G., Yakushev, N.N. The effectiveness of training with hypercapnic hypoxia in the rehabilitation of ischemic brain damage in the experiment. *Bulletin of Restorative Medicine*. 2008;2:59–61.
- Kulikov, V.P., Bespalov, A.G., Yakushev, N.N. The state of cerebral hemodynamics with longterm adaptation to hypercapnic hypoxia. *Russian Physiological Journal*. n. a. I. M. Sechenov. 2008;2:191–197.

- Kulikov, V.P., Osipov, I.S., Tregub, P.P. Optimal parameters of hypercapnic hypoxia effect for increasing resistance to acute hypoxia. *Aerospace and Ecology Medicine*. 2015;49(5):25–28.
- Kulikov, V.P., Tregub, P.P., Kovzelev, P.D., Dorokhov, E.A., Belousov, A.A. Hypercapnia — an alternative hypoxia signal stimulus for increasing HIF-1a and erythropoietin in the brain. *Patologicheskaya Fiziologiya i Eksperimental'naya Terapiya = Pathophysiology and Experimental Therapy*. 2015;3:34–37.
- Kuroda, J., Ago, T., Nishimura, A., Nakamura, K., Matsuo, R., Wakisaka, Y., Kamouchi, M., & Kitazono, T. (2014). Nox4 is a major source of superoxide production in human brain pericytes. *Journal of Vascular Research*, 51(6), 429–438.
- Kushwah, N., Jain, V., Deep, S., Prasad, D., Singh, S. B., & Khan, N. (2016). Neuroprotective role of intermittent hypobaric hypoxia in unpredictable chronic mild stress induced depression in rats. *PLoS ONE*, 11(2), e0149309.
- Kuznetsov, V.I., Solkin, A.A., Belyavsky, N.N. The effectiveness of the use of interval normobaric hypoxic therapy in patients with ischemic stroke in the recovery period. *Bulletin of Vitebsk State Medical University*. 2013;12(3):37–42.
- Lahiri, S., Forster, R.E., 2nd. CO<sub>2</sub>/H<sup>+</sup> sensing: Peripheral and central chemoreception. *Int. J. Biochem. Cell Biol.* 2003, 35, 1413–1435.
- Lam, K. K., Kunder, S., Wong, J., Doufas, A. G., & Chung, F. (2016). Obstructive sleep apnea, pain, and opioids: Is the riddle solved? *Current Opinion in Anaesthesiology*, 29(1), 134–140.
- Lasitsa, T.S., Morozova, N.A. The rationale for the use of a hypercapnic mixture in the aerosol therapy of patients with diseases of the lungs. *Probl Tuberk*. 1990;5:44–48.
- Latshang, T. D., Tardent, R. P. M., Furian, M., Flueck, D., Segitz, S. D., Mueller-Mottet, S., Kohler, M., Ulrich, S., & Bloch, K. E. (2019). Sleep and breathing disturbances in patients with chronic obstructive pulmonary disease traveling to altitude: A randomized trial. *Sleep*, 42(1). <https://doi.org/10.1093/sleep/zsy203>.
- Lavie, L. (2005). Sleep-disordered breathing and cerebrovascular disease: A mechanistic approach. *Neurologic Clinics*, 23(4), 1059–1075.
- Lavie, P., & Lavie, L. (2009). Unexpected survival advantage in elderly people with moderate sleep apnoea. *Journal of Sleep Research*, 18(4), 397–403.
- Lee, D. S., Badr, M. S., & Mateika, J. H. (2009). Progressive augmentation and ventilatory long-term facilitation are enhanced in sleep apnoea patients and are mitigated by antioxidant administration. *The Journal of Physiology*, 587(22), 5451–5467.
- Lee, E. S., Smith, W. E.d, Quach, H. T., Jones, B. D., Santilli, S. M., & Vatassery, G. T. (2005). Moderate hyperoxia (40%) increases antioxidant levels in mouse tissue. *Journal of Surgical Research*, 127(2), 80–84.
- Lee, J., Kim, H. S., Kim, S. M., Kim, D. I., & Lee, C. W. (2018). Hypoxia upregulates mitotic cyclins which contribute to the multipotency of human mesenchymal stem cells by expanding proliferation lifespan. *Molecules and Cells*, 41(3), 207–213.
- Lee, P., Chandel, N. S., & Simon, M. C. (2020). Cellular adaptation to hypoxia through hypoxia inducible factors and beyond. *Nature Reviews Molecular Cell Biology*, 21(5), 268–283.
- Lee, W.-C., Guntur, A. R., Long, F., & Rosen, C. J. (2017). Energy metabolism of the osteoblast: Implications for osteoporosis. *Endocrine Reviews*, 38(3), 255–266.
- Lendahl, U., Lee, K. L., Yang, H., & Poellinger, L. (2009). Generating specificity and diversity in the transcriptional response to hypoxia. *Nature Reviews Genetics*, 10(12), 821–832.
- Lestón Pinilla, L., Ugun-Klusek, A., Rutella, S., & De Girolamo, L. A. (2021). Hypoxia signaling in Parkinson's Disease: There is use in asking "What HIF?". *Biology*, 10, 723.
- Li, G., Guan, Y., Gu, Y., Guo, M., Ma, W., Shao, Q., Liu, J., & Ji, X. (2023). Intermittent hypoxic conditioning restores neurological dysfunction of mice induced by long-term hypoxia. *CNS Neuroscience & Therapeutics*, 29(1), 202–215.
- Li, S., Hafeez, A., Noorulla, F., Geng, X., Shao, G., Ren, C., Lu, G., Zhao, H., Ding, Y., & Ji, X. (2017). Preconditioning in neuroprotection: From hypoxia to ischemia. *Progress in Neurobiology*, 157, 79–91.
- Li, X.-L., Wang, W.-G., Li, M.-X., Liu, T.-L., Tian, X.-Y.u, & Wu, L. (2022). Effects of altitude and duration of differing levels of hypoxic exposure on hypoxia-inducible factor-1α in rat tissues. *High Altitude Medicine & Biology*, 23(2), 173–184.
- Li, Y., Zhang, Y., & Zhang, Y. (2018). Research advances in pathogenesis and prophylactic measures of acute high altitude illness. *Respiratory Medicine*, 145, 145–152.
- Lindholm, M. E., & Rundqvist, H. (2016). Skeletal muscle hypoxia-inducible factor-1 and exercise. *Experimental Physiology*, 101(1), 28–32.
- Linthwaite, V.L., Pawloski, W., Pegg, H.B., Townsend, P.D., Thomas, M.J., So, V.K.H., Brown, A.P., Hodgson, D.R.W., Lorimer, G.H., Fushman, D., et al. Ubiquitin is a carbon dioxide-binding protein. *Sci. Adv.* 2021, 7, eabi5507.
- Liu, L., Cash, T. P., Jones, R. G., Keith, B., Thompson, C. B., & Simon, M. C. (2006). Hypoxia-induced energy stress regulates mRNA translation and cell growth. *Molecular cell*, 21(4), 521–531.
- Liu, X., & Hajnoczky, G. (2011). Altered fusion dynamics underlie unique morphological changes in mitochondria during hypoxia-reoxygenation stress. *Cell Death & Differentiation*, 18, 1561–1572.
- Lizamore, C. A., & Hamlin, M. J. (2017). The use of simulated altitude techniques for beneficial cardiovascular health outcomes in nonathletic, sedentary, and clinical populations: A literature review. *High Altitude Medicine & Biology*, 18(4), 305–321.
- López-Barneo, J., González-Rodríguez, P., Gao, L., Fernández-Agüera, M. C., Pardal, R., & Ortega-Sáenz, P. (2016). Oxygen sensing by the carotid body: Mechanisms and role in adaptation to hypoxia. *American Journal of Physiology. Cell Physiology*, 310(8), C629–C642.

- Luks, A. M., & Hackett, P. H. (2022). Medical conditions and high-altitude travel. *New England Journal of Medicine*, 386(4), 364–373.
- Lukyanova L.D. (2011) Modern problems of adaptation to hypoxia. Signaling mechanisms and their role in systemic regulation. *Patol Fiziol Eksp Ter* 1:3–19.
- Lukyanova L.D., Germanova E.L., Kopaladze R.A. (2009) Development of resistance of an organism under various conditions of hypoxic preconditioning: role of the hypoxic period and reoxygenation. *Bull Exp Biol Med* 147(4):400–404.
- Lyamina, N. P., Lyamina, S. V., Senchiknin, V. N., Mallet, R. T., Downey, H. F., & Manukhina, E. B. (2011). Normobaric hypoxia conditioning reduces blood pressure and normalizes nitric oxide synthesis in patients with arterial hypertension. *Journal of Hypertension*, 29(11), 2265–2272.
- Ma, M.-C., Qian, H., Ghassemi, F., Zhao, P., & Xia, Y. (2005). Oxygen-sensitive ( $\delta$ -opioid receptor-regulated survival and death signals: Novel insights into neuronal preconditioning and protection. *The Journal of Biological Chemistry*, 280(16), 16208–16218.
- Mackenzie, R., Maxwell, N., Castle, P., Brickley, G., & Watt, P. (2011). Acute hypoxia and exercise improve insulin sensitivity (S(I) (2\*)) in individuals with type 2 diabetes. *Diabetes Metabolism Research and Reviews*, 27(1), 94–101.
- Macura, M., Ban Frangez, H., Cankar, K., Finžgar, M., Frangez, I. The effect of transcutaneous application of gaseous CO<sub>2</sub> on diabetic chronic wound healing—A double-blind randomized clinical trial. *Int. Wound J.* 2020, 17, 1607–1614.
- Mallet, R. T., Burtscher, J., Richalet, J.-P., Millet, G. P., & Burtscher, M. (2021). Impact of high altitude on cardiovascular health: Current perspectives. *Vascular Health and Risk Management*, 17, 317–335.
- Mallet, R. T., Ryoo, M.-G., Williams, A. G., Howard, L., & Downey, H. F. (2006). Beta1-Adrenergic receptor antagonism abrogates cardioprotective effects of intermittent hypoxia. *Basic Research in Cardiology*, 101(5), 436–446.
- Malte, H., Lykkeboe, G. The bohr/haldane effect: A model-based uncovering of the full extent of its impact on O<sub>2</sub> delivery to and CO<sub>2</sub> removal from tissues. *J. Appl. Physiol.* 2018, 125, 916–922.
- Manella, G., Aviram, R., Bolshette, N., Muvkadi, S., Golik, M., Smith, D. F., & Asher, G. (2020). Hypoxia induces a time- and tissue-specific response that elicits intertissue circadian clock misalignment. *Proceedings of the National Academy of Sciences USA*, 117(1), 779–786.
- Manimmanakorn, A., Manimmanakorn, N., Taylor, R., Draper, N., Billaut, F., Shearman, J. P., & Hamlin, M. J. (2013). Effects of resistance training combined with vascular occlusion or hypoxia on neuromuscular function in athletes. *European Journal of Applied Physiology*, 113(7), 1767–1774.
- Manukhina, E. B., Belkina, L. M., Terekhina, O. L., Abramochkin, D. V., Smirnova, E. A., Budanova, O. P., Mallet, R. T., & Downey, H. F. (2013). Normobaric, intermittent hypoxia conditioning is cardio- and vasoprotective in rats. *Experimental Biology and Medicine*, 238(12), 1413–1420.
- Manukhina, E. B., Downey, H. F., & Mallet, R. T. (2006). Role of nitric oxide in cardiovascular adaptation to intermittent hypoxia. *Experimental Biology and Medicine*, 231(4), 343–365.
- Manukhina, E. B., Downey, H. F., Shi, X., & Mallet, R. T. (2016). Intermittent hypoxia training protects cerebrovascular function in Alzheimer's disease. *Experimental Biology and Medicine*, 241(12), 1351–1363.
- Manukhina, E. B., Pshennikova, M. G., Goryacheva, A. V., Khomenko, I. P., Mashina, S. Yu., Pokidyshev, D. A., & Malyshev, I. Y. (2008). Role of nitric oxide in prevention of cognitive disorders in neurodegenerative brain injuries in rats. *Bulletin of Experimental Biology and Medicine*, 146(4), 391–395.
- Manukhina, E. B., Tseilikman, V. E., Karpenko, M. N., Pestereva, N. S., Tseilikman, O. B., Komelkova, M. V., Kondashevskaya, M. V., Goryacheva, A. V., Lapshin, M. S., Platkovskii, P. O., Sarapultsev, A. P., Alliluev, A. V., & Downey, H. F. (2020). Intermittent hypoxic conditioning alleviates post-traumatic stress disorder-induced damage and dysfunction of rat visceral organs and brain. *International Journal of Molecular Sciences*, 21(1), 345.
- Manukhina, E. B., Tseilikman, V. E., Tseilikman, O. B., Komelkova, M. V., Kondashevskaya, M. V., Goryacheva, A. V., Lapshin, M. S., Platkovskii, P. O., Alliluev, A. V., & Downey, H. F. (2018). Intermittent hypoxia improves behavioral and adrenal gland dysfunction induced by post-traumatic stress disorder in rats. Maruyama, K.; Okada, T.; Ueha, T.; Isohashi, K.; Ikeda, H.; Kanai, Y.; Sasaki, K.; Gentsu, T.; Ueshima, E.; Sofue, K.; et al. In vivo evaluation of percutaneous carbon dioxide treatment for improving intratumoral hypoxia using 18F-fluoromisonidazole PET-CT. *Oncol. Lett.* 2021, 21, 207. *Journal of Applied Physiology*, 125(3), 931–937.
- Maruyama, K., Okada, T., Ueha, T., Isohashi, K., Ikeda, H., Kanai, Y., Sasaki, K., Gentsu, T., Ueshima, E., Sofue, K., et al. In vivo evaluation of percutaneous carbon dioxide treatment for improving intratumoral hypoxia using 18F-fluoromisonidazole PET-CT. *Oncol. Lett.* 2021, 21, 207.
- Mashina, S. Y., Aleksandrin, V. V., Goryacheva, A. V., Vlasova, M. A., Vanin, A. F., Malyshev, I. Y., & Manukhina, E. B. (2006). Adaptation to hypoxia prevents disturbances in cerebral blood flow during neuro-degenerative process. *Bulletin of Experimental Biology and Medicine*, 142(2), 169–172.
- Mateika, J. H., Mendello, C., Obeid, D., & Badr, M. S. (2004). Peripheral chemoreflex responsiveness is increased at elevated levels of carbon dioxide after episodic hypoxia in awake humans. *Journal of Applied Physiology*, 96(3), 1197–1205. discussion 1196.
- Mateika, J. H., Panza, G., Alex, R., & El-Chami, M. (2018). The impact of intermittent or sustained carbon dioxide on intermittent hypoxia initiated respiratory plasticity. What is the effect of these combined stimuli on apnea severity? *Respiratory Physiology & Neurobiology*, 256, 58–66.
- Matsumoto, T., Tanaka, M., Ikeji, T., Maeshige, N., Sakai, Y., Akisue, T., Kondo, H., Ishihara, A., Fujino, H. Application of transcutaneous carbon dioxide improves capillary regression of skeletal muscle in hyperglycemia. *J. Physiol. Sci.* 2019, 69, 317–326.

- Matsumoto, T., Tanaka, M., Nakanish, R., Takuwa, M., Hirabayashi, T., Ono, K., Ikeji, T., Maeshige, N., Sakai, Y., Akisue, T., et al. Transcutaneous carbon dioxide attenuates impaired oxidative capacity in skeletal muscle in hyperglycemia model. *Gen. Physiol. Biophys.* 2019, 38, 237–244.
- Matu, J., Gonzalez, J. T., Ispoglou, T., Duckworth, L., & Deighton, K. (2018). The effects of hypoxia on hunger perceptions, appetite-related hormone concentrations and energy intake: A systematic review and meta-analysis. *Appetite*, 125, 98–108.
- Maulik, N., Engelman, R.M., Rouson, J.A. et al. Ischemic preconditioning reduces apoptosis by up regulating Anti-death gene Bcl-2 // *Circulation*, 1999. -v.100 (Suppl 2). -P.369-375.
- McLaughlin B., Hartnett K.A., Erhardt J.A. et al (2003) Caspase 3 activation is essential for neuroprotection in preconditioning. *Proc Natl Acad Sci USA* 100:715–720.
- Meerson, F.Z., Malyshev, I.Yu. Adaptation to stress increases the heart resistance to ischemic and reperfusion arrhythmias. *J Mol Cell Cardiol.* 1989;21(3):299–303.
- Menuet, C., Connelly, A. A., Bassi, J. K., Melo, M. R., Le, S., Kamar, J., Kumar, N. N., McDougall, S. J., McMullan, S., & Allen, A. M. (2020). PreBötzinger complex neurons drive respiratory modulation of blood pressure and heart rate. *eLife*, 9, e57288.
- Metallo, C. M., Gameiro, P. A., Bell, E. L., Mattaini, K. R., Yang, J., Hiller, K., Jewell, C. M., Johnson, Z. R., Irvine, D. J., Guarante, L., Kelleher, J. K., Vander Heiden, M. G., Iliopoulos, O., & Stephanopoulos, G. (2011). Reductive glutamine metabolism by IDH1 mediates lipogenesis under hypoxia. *Nature*, 481(7381), 380–384.
- Meyer, M. J., Mordukhovich, I., Wollenius, G. A., Mittleman, M. A., McCracken, J. P., Coull, B. A., & Mcneely, E. (2019). Changes in heart rate and rhythm during a crossover study of simulated commercial flight in older and vulnerable participants. *Frontiers in Physiology*, 10, 1339.
- Midha, A. D., Zhou, Y., Queliconi, B. B., Barrios, A. M., Haribowo, A. G., Chew, B. T. L., Fong, C. O. Y., Blecha, J. E., Vanbrocklin, H., Seo, Y., & Jain, I. H. (2023). Organ-specific fuel rewiring in acute and chronic hypoxia redistributes glucose and fatty acid metabolism. *Cell Metabolism*, 35(3), 504–516.e5. e505.
- Milano, G., Abruzzo, P. M., Bolotta, A., Marini, M., Terraneo, L., Ravara, B., Gorza, L., Vitadello, M., Burattini, S., Curzi, D., Falcieri, E., Von Segesser, L. K., & Samaja, M. (2013). Impact of the phosphatidylinositol 3-kinase signaling pathway on the cardioprotection induced by intermittent hypoxia. *PLoS ONE*, 8(10), e76659.
- Miller, A. J., Sauder, C. L., Cauffman, A. E., Blaha, C. A., & Leuenberger, U. A. (2017). Endurance training attenuates the increase in peripheral chemoreflex sensitivity with intermittent hypoxia. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 312(2), R223–R228.
- Millet, G. P., Roels, B., Schmitt, L., Woorons, X., & Richalet, J. P. (2010). Combining hypoxic methods for peak performance. *Sports Medicine*, 40(1), 1–25.
- Minamiyama, M., Yamamoto, A. Direct evidence of the vasodilator action of carbon dioxide on subcutaneous microvasculature in rats by use of intra-vital video-microscopy. *J. Biorheol.* 2010, 24, 42–46.
- Mole, D. R., Blancher, C., Copley, R. R., Pollard, P. J., Gleadle, J. M., Ragoussis, J., & Ratcliffe, P. J. (2009). Genome-wide association of hypoxia-inducible factor (HIF)-1 $\alpha$  and HIF-2 $\alpha$  DNA binding with expression profiling of hypoxia-inducible transcripts. *Journal of Biological Chemistry*, 284(25), 16767–16775.
- Montalto, A.S., Currò, M., Russo, T., Visalli, G., Impellizzeri, P., Antonuccio, P., Arena, S., Borruto, F.A., Scalfari, G., Lentile, R., et al. In Vitro CO<sub>2</sub>-induced ROS production impairs cell cycle in SH-SY5Y neuroblastoma cells. *Pediatr. Surg. Int.* 2013, 29, 51–59.
- Morelli, C., Badr, M. S., & Mateika, J. H. (2004). Ventilatory responses to carbon dioxide at low and high levels of oxygen are elevated after episodic hypoxia in men compared with women. *Journal of Applied Physiology*, 97(5), 1673–1680.
- Moreno-Domínguez, A., Ortega-Sáenz, P., Gao, L., Colinas, O., García-Flores, P., Bonilla-Henao, V., Aragónés, J., Hüttemann, M., Grossman, L. I., Weissmann, N., Sommer, N., & López-Barneo, J. (2020). Acute O<sub>2</sub> sensing through HIF2 $\alpha$ -dependent expression of atypical cytochrome oxidase subunits in arterial chemoreceptors. *Science Signaling*, 13(615), eaay9452.
- Morishima, T., Hasegawa, Y., Sasaki, H. Effects of different periods of hypoxic training on glucose metabolism and insulin sensitivity. *Clin Physiol Funct Imaging*. 2014. doi:10.1111/cpf.12133.
- Moulin, S., Arnaud, C., Bouyon, S., Pépin, J.-L., Godin-Ribuot, D., & Belaïdi, E. (2020). Curcumin prevents chronic intermittent hypoxia-induced myocardial injury. *Therapeutic Advances in Chronic Disease*, 11, 204062232092210.
- Muharlyamov, F.Yu., Smirnova, M.I., Bedritsky, S.A. Interval hypoxic training for arterial hypertension. *Questions of Balneology, Physiotherapy and Physical Therapy*. 2006;2:5–6.
- Muangritdech, N., Hamlin, M. J., Sawanyawisuth, K., Prajumwongs, P., Saengjan, W., Wonnabussapawich, P., Manimmanakorn, N., & Manimmanakorn, A. (2020). Hypoxic training improves blood pressure, nitric oxide and hypoxia-inducible factor-1 alpha in hypertensive patients. *European Journal of Applied Physiology*, 120(8), 1815–1826.
- Mulkey, D. K., Rosin, D. L., West, G., Takakura, A. C., Moreira, T. S., Bayliss, D. A., & Guyenet, P. G. (2007). Serotonergic neurons activate chemosensitive retrotrapezoid nucleus neurons by a pH-independent mechanism. *Journal of Neuroscience*, 27(51), 14128–14138.
- Murphy, B.J. Regulation of malignant progression by the hypoxia-sensitive transcription factors HIF-1 $\alpha$  and MTF-1. *Comp Biochem Physiol B Biochem Mol Biol*. 2004;139(3):495–507.
- Nanduri, J., Vaddi, D. R., Khan, S. A., Wang, N., Makarenko, V., & Prabhakar, N. R. (2013). Xanthine oxidase mediates hypoxia-inducible factor-2 $\alpha$  degradation by intermittent hypoxia. *PLoS ONE*, 8(10), e75838.
- Navarrete-Opazo, A., & Mitchell, G. S. (2014). Therapeutic potential of intermittent hypoxia: A matter of dose. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 307(10), R1181–R1197.
- Neckár J., Papousek F., Novákóvá O. et al (2002) Cardioprotective effects of chronic hypoxia and ischaemic preconditioning are not additive. *Basic Res Cardiol* 97(2):161–167.

- Nees, S., Gerlach, E. Adenine Nucleotide and Adenosine Metabolism in Cultured Coronary Endothelial Cells: Formation and Release of Adenine Compounds and Possible Functional Implications. In *Regulatory Function of Adenosine*; Berne, R.M., Rall, T.W., Rubio, R., Eds.; Springer: Boston, MA, USA, 1983; pp. 347–360.
- Németh, B., Kiss, I., Ajtay, B., Péter, I., Kreska, Z., Cziráki, A., Horváth, I.G., Ajtay, Z. Transcutaneous carbon dioxide treatment is capable of reducing peripheral vascular resistance in hypertensive patients. *Vivo* 2018, 32, 1555–1559.
- Nemeth, B., Kiss, I.; Jencsik, T., Peter, I., Kreska, Z., Koszegi, T., Miseta, A., Kustan, P., Boncz, I., Laczo, A., et al. Angiotensin-converting enzyme inhibition improves the effectiveness of transcutaneous carbon dioxide treatment. *vivo* 2017, 31, 425–428.
- Netzer, N., Gatterer, H., Faulhaber, M., Burtscher, M., Pramsohler, S., & Pesta, D. (2015). Hypoxia, oxidative stress and fat. *Biomolecules*, 5(2), 1143–1150.
- Nichols, N. L., & Mitchell, G. S. (2021). Mechanisms of severe acute intermittent hypoxia-induced phrenic long-term facilitation. *Journal of Neurophysiology*, 125(4), 1146–1156.
- Niikura, T., Iwakura, T., Omori, T., Lee, S.Y., Sakai, Y., Akisue, T., Oe, K., Fukui, T., Matsushita, T., Matsumoto, T., et al. Topical cutaneous application of carbon dioxide via a hydrogel for improved fracture repair: Results of phase I clinical safety trial. *BMC Musculoskelet. Disord.* 2019, 20, 563.
- Nishimoto, H., Inui, A., Ueha, T., Inoue, M., Akahane, S., Harada, R., Mifune, Y., Kokubu, T., Nishida, K., Kuroda, R., et al. Transcutaneous carbon dioxide application with hydrogel prevents muscle atrophy in a rat sciatic nerve crush model. *J. Orthop. Res.* 2018, 36, 1653–1658.
- Obrenovich T.P. (2008) Molecular physiology of preconditioning-induced brain tolerance to ischemia. *Physiol Rev* 88:211–247.
- Oda, T., Iwakura, T., Fukui, T., Oe, K., Mifune, Y., Hayashi, S., Matsumoto, T., Matsushita, T., Kawamoto, T., Sakai, Y., et al. Effects of the duration of transcutaneous CO<sub>2</sub> application on the facilitatory effect in rat fracture repair. *J. Orthop. Sci.* 2020, 25, 886–891.
- Oe, K., Ueha, T., Sakai, Y., Nikura, T., Lee, S.Y., Koh, A., Hasegawa, T., Tanaka, M., Miwa, M., Kurosaka, M. The effect of transcutaneous application of carbon dioxide (CO<sub>2</sub>) on skeletal muscle. *Biochem. Biophys. Res. Commun.* 2011, 407, 148–152.
- Ogoh, S., Nagaoka, R., Mizuno, T., Kimura, S., Shidahara, Y., Ishii, T., Kudoh, M., Iwamoto, E. Acute vascular effects of carbonated warm water lower leg immersion in healthy young adults. *Physiol. Rep.* 2016, 4, e13046.
- Olcina, M. M., Foskolou, I. P., Anbalagan, S., Senra, J. M., Pires, I. M., Jiang, Y., Ryan, A. J., & Hammond, E. M. (2013). Replication stress and chromatin context link ATM activation to a role in DNA replication. *Molecular cell*, 52(5), 758–766.
- Onishi, Y., Akisue, T., Kawamoto, T., Ueha, T., Hara, H., Toda, M., Harada, R., Minoda, M., Morishita, M., Sasaki, R., et al. Transcutaneous application of CO<sub>2</sub> enhances the antitumor effect of radiation therapy in human malignant fibrous histiocytoma. *Int. J. Oncol.* 2014, 45, 732–738.
- Onishi, Y., Kawamoto, T. Transcutaneous application of carbon dioxide (CO<sub>2</sub>) enhances chemosensitivity by reducing hypoxic conditions in human malignant fibrous histiocytoma. *J. Cancer Sci. Ther.* 2012, 4, 74–181.
- Ostadal, B., Kolar, F. Cardiac adaptation to chronic high-altitude hypoxia: beneficial and adverse effects. *Respir Physiol Neurobiol.* 2007;158(2–3):224–236.
- Pamerter, M. E., & Powell, F. L. (2016). Time domains of the hypoxic ventilatory response and their molecular basis. *Comprehensive Physiology*, 6(3), 1345–1385.
- Panza, G. S., Alex, R. M., Yokhana, S. S., Lee Pioszak, D. S., Badr, M. S., & Mateika, J. H. (2019). Increased oxidative stress, loop gain and the arousal threshold are clinical predictors of increased apnea severity following exposure to intermittent hypoxia. *Nature and Science of Sleep*, 11, 265–279.
- Panza, G. S., Burtscher, J., & Zhao, F. (2023). Intermittent Hypoxia: A call for harmonization in terminology. *Journal of Applied Physiology*, 135(4), 886–890.
- Panza, G. S., Puri, S., Lin, H.-S., Badr, M. S., & Mateika, J. H. (2022). Daily exposure to mild intermittent hypoxia reduces blood pressure in male OSA patients with hypertension. *American Journal of Respiratory and Critical Care Medicine*, 205(8), 949–958.
- Panza, G. S., Puri, S., Lin, H.-S., & Mateika, J. H. (2022). Divergent ventilatory and blood pressure responses are evident following repeated daily exposure to mild intermittent hypoxia in males with OSA and hypertension. *Frontiers in Physiology*, 13, 897978.
- Papandreou, I., Cairns, R. A., Fontana, L., Lim, A.i L., & Denko, N. C. (2006). HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metabolism*, 3(3), 187–197.
- Parati, G., Ochoa, J. E., Torlasco, C., Salvi, P., Lombardi, C., & Bilo, G. (2015). Aging, high altitude, and blood pressure: A complex relationship. *High Altitude Medicine & Biology*, 16(2), 97–109.
- Pechkina, K.G., Kulikov, V.P., Shcherbakov, P.L., Lobanov, Yu.F. Treatment of chronic erosive gastroduodenitis in children with hypercapnic hypoxia. *Eksperimental'naya i Klinicheskaya Gastroenterologiya = Experimental and Clinical Gastroenterology*. 2011;65(1):28–30.
- Pernett, F., Schagatay, F., Vildevi, C., & Schagatay, E. (2021). Spleen contraction during sudden eupneic hypoxia elevates hemoglobin concentration. *Frontiers in Physiology*, 12, 729123.
- Pesta, D., Heieis, J., Hand, O., Frings-Meuthen, P., Marcus, K., Clemen, C. S., Levine, B., Sadek, H., Hoffmann, F., Limper, U., Jordan, J., Sies, W., Tank, J., Zange, J., & Rittweger, J. (2023). Case report: Muscle wasting during severe sustained hypoxia in two professional mountaineers. *Medicine and Science in Sports and Exercise*, 55(3), 335–341.
- Phillis, J.W., O'Regan, M.H., Song, D. Further evidence for the role of adenosine in hypercapnia/acidosis-evoked coronary flow regulation. *Gen. Pharmacol.* 1999, 33, 431–437.
- Potievskaya, V.I. Mechanisms of therapeutic and preventive effects of hypoxia in hypertension. *Journal of Physiology*. 1993;39 (2):94–107.

- Prasad, B., Morgan, B. J., Gupta, A., Pegelow, D. F., Teodorescu, M., Dopp, J. M., & Dempsey, J. A. (2020). The need for specificity in quantifying neurocirculatory vs. respiratory effects of eucapnic hypoxia and transient hyperoxia. *The Journal of Physiology*, 598(21), 4803–4819.
- Pueyo, M. E., Gonzalez, W., Nicoletti, A., Savoie, F., Arnal, J.-F., & Michel, J.-B. (2000). Angiotensin II stimulates endothelial vascular cell adhesion molecule-1 via nuclear factor- $\kappa$ B activation induced by intracellular oxidative stress. *Arteriosclerosis, Thrombosis, and Vascular biology*, 20(3), 645–651.
- Pulido-Escribano, V., Torrecillas-Baena, B., Camacho-Cardenosa, M., Dorado, G., Gálvez-Moreno, M. Á., & Casado-Díaz, A. (2022). Role of hypoxia preconditioning in therapeutic potential of mesenchymal stem-cell-derived extracellular vesicles. *World Journal of Stem Cells*, 14(7), 453–472.
- Puri, S., El-Chami, M., Shaheen, D., Ivers, B., Panza, G. S., Badr, M. S., Lin, H.-S., & Mateika, J. H. (2020). Variations in loop gain and arousal threshold during NREM sleep are affected by time of day over a 24-hour period in participants with obstructive sleep apnea. *Journal of Applied Physiology*, 129(4), 800–809.
- Puri, S., Panza, G., & Mateika, J. H. (2021). A comprehensive review of respiratory, autonomic and cardiovascular responses to intermittent hypoxia in humans. *Experimental Neurology*, 341, 113709.
- Raberin, A., Burtscher, J., Connes, P., & Millet, G. P. (2022). Hypoxia and hemorheological properties in older individuals. *Ageing Research Reviews*, 79, 101650.
- Rachok, L.V., Dubovik, T.A., Bulgak, A.G. Use of intermittent normobaric hypoxotherapy in preoperative preparation for coronary bypass surgery in patients with ischemic cardiomyopathy. *Cardiology in Belarus*. 2011;17:28–45.
- Radermacher, K. A., Wingler, K., Langhauser, F., Altenhofer, S., Kleikers, P., Hermans, J. J., Hrabe de Angelis, M., Kleinschmitz, C., & Schmidt, H. H. (2013). Neuroprotection after stroke by targeting NOX4 as a source of oxidative stress. *Antioxidants & Redox Signaling*, 18(12), 1418–1427.
- Rafibekova, I.S., Dzhumangulova, A.S., Usubaliev, N.N. Treatment of hypertension with hypobaric pressure chamber hypoxia and mid-mountain climate. XIX All-Union Congress of Physicians. Tashkent. 1987;3:29–30.
- Rahimi, N. The ubiquitin-proteasome system meets angiogenesis. *Mol. Cancer Res.* 2012, 11, 538–548.
- Rasbach, K. A., Gupta, R. K., Ruas, J. L., Wu, J., Naseri, E., Estall, J. L., & Spiegelman, B. M. (2010). PGC-1alpha regulates a HIF2alpha-dependent switch in skeletal muscle fiber types. *Proceedings of the National Academy of Sciences, USA*, 107(50), 21866–21871.
- Reeves, J.T., Weil, J.V. Chronic mountain sickness. A view from the crow's nest. *Adv Exp Med Biol.* 2001;502:419–437.
- Reglin, B., Pries, A.R. Metabolic control of microvascular networks: Oxygen sensing and beyond. *J. Vasc. Res.* 2014, 51, 376–392.
- Richardson, M. X., Lodin, A., Reimers, J., & Schagatay, E. (2008). Short-term effects of normobaric hypoxia on the human spleen. *European Journal of Applied Physiology*, 104(2), 395–399.
- Ristow, M., & Schmeisser, K. (2014). Mitohormesis: Promoting health and lifespan by increased levels of reactive oxygen species (ROS). *Dose Response*, 12(2), 288–341.
- Rodríguez, C., Muñoz, M., Contreras, C., Prieto, D. AMPK, metabolism, and vascular function. *FEBS J.* 2021, 288, 3746–3771.
- Rodriguez, F. A., Casas, H., Casas, M., Pagés, T., Rama, R., Ricart, A., Ventura, J. L., Ibáñez, J., & Viscor, G. (1999). Intermittent hypobaric hypoxia stimulates erythropoiesis and improves aerobic capacity. *Medicine & Science in Sports & Exercise*, 31(2), 264–268.
- Roy, J., Galano, J.-M., Durand, T., Le Guennec, J.-Y., & Chung-Yung Lee, J. (2017). Physiological role of reactive oxygen species as promoters of natural defenses. *FASEB Journal*, 31(9), 3729–3745.
- Ruiz L., Penalosa D. Altitude and hypertension. *Mayo Clin Proc.* 1977;52(7):442–445.
- Rybnikova, E., Mironova, V., Pivina, S., Tulkova, E., Ordyan, N., Nalivaeva, N., Turner, A., & Samoilov, M. (2007). Involvement of the hypothalamic-pituitary-adrenal axis in the antidepressant-like effects of mild hypoxic preconditioning in rats. *Psychoneuroendocrinology*, 32(7), 813–823.
- Rybnikova, E., Mironova, V., Pivina, S., Tulkova, E., Ordyan, N., Vataeva, L., Vershinina, E., Abritalin, E., Kolchev, A., Nalivaeva, N., Turner, A. J., & Samoilov, M. (2007). Antidepressant-like effects of mild hypoxia preconditioning in the learned helplessness model in rats. *Neuroscience Letters*, 417(3), 234–239.
- Rybnikova, E., & Nalivaeva, N. (2021). Glucocorticoid-dependent mechanisms of brain tolerance to hypoxia. *International Journal of Molecular Sciences*, 22(15), 7982.
- Rybnikova, E., & Samoilov, M. (2015). Current insights into the molecular mechanisms of hypoxic pre- and postconditioning using hypobaric hypoxia. *Frontiers in Neuroscience*, 9, 388.
- Rybnikova E., Sitnik N., Gluschenko T. et al (2006) The preconditioning modified neuronal expression of apoptosis-related proteins of Bcl-2 superfamily following severe hypobaric hypoxia in rats. *Brain Res* 1089(1):195–202. <https://doi.org/10.1016/j.brainres.2006.03.053>.
- Rybnikova, E. A., Mironova, V. I., Pivina, S. G., Ordyan, N. E., Tulkova, E. I., & Samoilov, M. O. (2008). Hormonal mechanisms of neuroprotective effects of the mild hypoxic preconditioning in rats. *Doklady Biological Sciences*, 421(1), 239–240.
- Rybnikova, E. A., Nalivaeva, N. N., Zenko, M. Y., & Baranova, K. A. (2022). Intermittent hypoxic training as an effective tool for increasing the adaptive potential, endurance and working capacity of the brain. *Frontiers in Neuroscience*, 16, 941740.
- Rybnikova, E. A., Samoilov, M. O., Mironova, V. I., Tyul'kova, E. I., Pivina, S. G., Vataeva, L. A., Ordyan, N. É., Abritalin, E. Y.u., & Kolchev, A. I. (2008). The possible use of hypoxic preconditioning for the prophylaxis of post-stress depressive episodes. *Neuroscience and Behavioral Physiology*, 38(7), 721–726.

- Ryou, M.-G., Chen, X., Cai, M., Wang, H., Jung, M. E., Metzger, D. B., Mallet, R. T., & Shi, X. (2021). Intermittent hypoxia training prevents deficient learning-memory behavior in mice modeling alzheimer's disease: A pilot study. *Frontiers in Aging Neuroscience*, 13, 674688.
- Ryou, M.-G., Sun, J., Oguayo, K. N., Manukhina, E. B., Downey, H. F., & Mallet, R. T. (2008). Hypoxic conditioning suppresses nitric oxide production upon myocardial reperfusion. *Experimental Biology and Medicine*, 233(6), 766–774.
- Saito, I., Hasegawa, T., Ueha, T., Takeda, D., Iwata, E., Arimoto, S., Sakakibara, A., Sakakibara, S., Akashi, M., Sakai, Y., et al. Effect of local application of transcutaneous carbon dioxide on survival of random-pattern skin flaps. *J. Plast. Reconstr. Aesthet. Surg.* 2018, 71, 1644–1651.
- Sakai, Y., Miwa, M., Oe, K., Ueha, T., Koh, A., Niikura, T., Iwakura, T., Lee, S.Y., Tanaka, M., Kurosaka, M. A novel system for transcutaneous application of carbon dioxide causing an "artificial bohr effect" in the human body. *PLoS ONE* 2011, 6, e24137.
- Sakai, Y., Ueha, T., Akisue, T., et al. Optimization of antitumor treatment conditions for transcutaneous CO<sub>2</sub> application: An *in vivo* study.(carbon dioxide ). *Onc Reports* 2017;37:3688.
- Sandbichler, A. M., Jansen, B., Peer, B. A., Paulitsch, M., Pelster, B., & Egg, M. (2018). Metabolic plasticity enables circadian adaptation to acute hypoxia in zebrafish cells. *Cellular Physiology and Biochemistry*, 46(3), 1159–1174.
- Savin, E., Bailliart, O., Bonnin, P., et al. Vasomotor Effects of Transcutaneous CO<sub>2</sub> in Stage II Peripheral Occlusive Arterial Disease. *Angiology* 1995;46:785-791.
- Sazontova, T. G., & Arkhipenko, I. V. (2005). [The role of free radical processes and redox-signalization in adaptation of the organism to changes in oxygen level]. *Rossiiiskii fiziologicheskii zhurnal imeni IM Sechenova*, 91(6), 636– 655.
- Sazontova, T.G., Zhukova, A.G., Anchishkina, N.A., Arkhipenko, Yu.V. HIF-1α Transcription Factor and Urgent Response Proteins in Acute Hypoxia and Adaptation. Abstracts of the 5th International Congress of Pathophysiology, Beijing, China. 2006. June 28-July 1 // *Chinese Journal of Pathophysiology*, 2006. -v.22. -№ 3 (Suppl.). -P.1-14.
- Schito, L., & Semenza, G. L. (2016). Hypoxia-inducible factors: Master regulators of cancer progression. *Trends in Cancer*, 2(12), 758–770.
- Schlittler, M., Gatterer, H., Turner, R., Regli, I. B., Woyke, S., Strapazzon, G., Rasmussen, P., Kob, M., Mueller, T., Goetze, J. P., Maillard, M., Van Hall, G., Feraille, E., & Siebenmann, C. (2021). Regulation of plasma volume in male lowlanders during 4 days of exposure to hypobaric hypoxia equivalent to 3500 m altitude. *The Journal of Physiology*, 599(4), 1083–1096.
- Schmidt, W. (2002). Effects of intermittent exposure to high altitude on blood volume and erythropoietic activity. *High Altitude Medicine & Biology*, 3(2), 167–176.
- Scholz, C. C., Cavadas, M. A. S., Tambuwala, M. M., Hams, E., Rodríguez, J., Kriegsheim, A. V., Cotter, P., Bruning, U., Fallon, P. G., Cheong, A., Cummins, E. P., & Taylor, C. T. (2013). Regulation of IL-1β-induced NF-κB by hydroxylases links key hypoxic and inflammatory signaling pathways. *Proceedings of the National Academy of Sciences, USA*, 110(46), 18490–18495.
- Schoos, A., Gabriel, C., Knab, V. M., & Fux, D. A. (2019). Activation of HIF-1α by δ-opioid receptors induces COX-2 expression in breast cancer cells and leads to paracrine activation of vascular endothelial cells. *The Journal of Pharmacology and Experimental Therapeutics*, 370(3), 480–489.
- Schuler, B., Vogel, J., Grenacher, B., Jacobs, R. A., Arras, M., & Gassmann, M. (2012). Acute and chronic elevation of erythropoietin in the brain improves exercise performance in mice without inducing erythropoiesis. *FASEB Journal*, 26(9), 3884–3890.
- Scott, B. R., Slattery, K. M., Sculley, D. V., Lockhart, C., & Dascombe, B. J. (2017). Acute physiological responses to moderate-load resistance exercise in hypoxia. *The Journal of Strength & Conditioning Research*, 31, 1973–1981.
- Selfridge, A.C., Cavadas, M.A.S., Scholz, C.C., et al. Hypercapnia Suppresses the HIF-dependent Adaptive Response to Hypoxia. *J Biol Chem* 2016;291:11800-11808.
- Semenza, G.L. Regulation of mammalian O<sub>2</sub> homeostasis by hypoxia-inducible factor 1. *Annu Rev Cell Dev Biol*. 1999;15:551–578. doi:10.1146/annurev.cellbio.15.1.551.
- Semenza, G.L. (2001). HIF-1 and mechanisms of hypoxia sensing. *Current Opinion in Cell Biology*, 13(2), 167– 171.
- Semenza, G.L. (2012). Hypoxia-inducible factors in physiology and medicine. *Cell*, 148(3), 399–408.
- Semenza, G.L. HIF-1 mediates metabolic responses to intratumoral hypoxia and oncogenic mutations. *J. Clin. Investigig.* 2013, 123, 3664–3671.
- Serebrovskaya T.V. Intermittent hypoxia research in the former Soviet Union and the commonwealth of independent states (CIS): history and review of the concept and selected applications. *High Alt Med Biol*. 2002;3(2):205–221.
- Serebrovska, T. V., Portnychenko, A. G., Drevytska, T. I., Portnychenko, V. I., Xi, L., Egorov, E., Gavalko, A. V., Naskalova, S., Chizhova, V., & Shatylo, V. B. (2017). Intermittent hypoxia training in prediabetes patients: Beneficial effects on glucose homeostasis, hypoxia tolerance and gene expression. *Experimental Biology and Medicine*, 242(15), 1542–1552.
- Serebrovskaya, T.V., Shatilo, V.B. Experience of using interval hypoxia for the prevention and treatment of diseases of the cardiovascular system. Overview. *Blood Circulation and Hemostasis*. 2014;(1-2):16-33.
- Serebrovska, Z. O., Serebrovska, T. V., Kholin, V. A., Tumanovska, L. V., Shysh, A. M., Pashevin, D. A., Goncharov, S. V., Stroy, D., Grib, O. N., Shatylo, V. B., Bachinskaya, N. Yu., Egorov, E., Xi, L., & Dosenko, V. E. (2019). Intermittent hypoxia-hyperoxia training improves cognitive function and decreases circulating biomarkers of Alzheimer's disease in patients with mild cognitive impairment: A pilot study. *International Journal of Molecular Sciences*, 20(21), 5405.

- Serebrovska, Z. O., Xi, L., Tumanovska, L. V., Shysh, A. M., Goncharov, S. V., Khetseriani, M., Kozak, T. O., Pasheviv, D. A., Dosenko, V. E., Virko, S. V., Kholin, V. A., Grib, O. N., Utko, N. A., Egorov, E., Polischuk, A. O., & Serebrovska, T. V. (2022). Response of circulating inflammatory markers to intermittent hypoxia-hyperoxia training in healthy elderly people and patients with mild cognitive impairment. *Life*, 12(3), 432.
- Serebrovskaya, T. V., Manukhina, E. B., Smith, M. L., Downey, H. F., & Mallet, R. T. (2008). Intermittent hypoxia: Cause of or therapy for systemic hypertension? *Experimental Biology and Medicine*, 233(6), 627–650.
- Serebrovskaya, T. V., Nikolsky, I. S., Nikolska, V. V., Mallet, R. T., & Ishchuk, V. A. (2011). Intermittent hypoxia mobilizes hematopoietic progenitors and augments cellular and humoral elements of innate immunity in adult men. *High Altitude Medicine & Biology*, 12(3), 243–252.
- Shatilo, V. B., Korkushko, O. V., Ischuk, V. A., Downey, H. F., & Serebrovskaya, T. V. (2008). Effects of intermittent hypoxia training on exercise performance, hemodynamics, and ventilation in healthy senior men. *High Altitude Medicine & Biology*, 9(1), 43–52.
- Sharp F.R., Ran R., Lu A. et al (2004) Hypoxic preconditioning protects against ischemic brain injury. *NeuroRx* 1:26–35.
- Shatilo V.B., Korkushko O.V., Ischuk V.A. et al (2008) Effects of intermittent hypoxia training on exercise performance, hemodynamics, and ventilation in healthy senior men. *High Altern Med Biol* 9(1):43–52.
- Sheikhbahaei, S. (2020). Physiology: New Insights into central oxygen sensing. *Current Biology*, 30(17), R1004–R1006.
- Shevchenko, Yu.L. (ed.) Hypoxia: adaptation, pathogenesis, clinic. St Petersburg: Elbi-SPb, 2000. 384 p.
- Shigemura, M., Lecuona, E., Szajner, J.I. Effects of hypercapnia on the lung. *J Physiol* 2017;595:2431-2437.
- Sinclair, S.E., Kregenow, D.A., Lamm, W.J.E., Starr, I.R., Chi, E.Y., Hlastala, M.P. Hypercapnic Acidosis Is Protective in an In Vivo Model of Ventilator-induced Lung Injury. *Amer J Resp and Crit Care Med* 2002;166:403-408.
- Shoag, J., Arany, Z. Regulation of hypoxia-inducible genes by PGC-1α. *Arterioscl Throm Vas*. 2010, 30, 662–666.
- Shpilka, T., & Haynes, C. M. (2018). The mitochondrial UPR: Mechanisms, physiological functions and implications in ageing. *Nature Reviews Molecular Cell Biology*, 19(2), 109–120.
- Siebenmann, C., Robach, P., & Lundby, C. (2017). Regulation of blood volume in lowlanders exposed to high altitude. *Journal of Applied Physiology*, 123(4), 957–966.
- Simon, M. C., & Keith, B. (2008). The role of oxygen availability in embryonic development and stem cell function. *Nature Reviews Molecular Cell Biology*, 9(4), 285–296.
- Simonenko, V.B., Ermolaev, A.L., Potievskaya, V.I., Stepaniants, O.S. Hypoxic therapy of arterial hypertension in patients with different blood pressure variability. *Klin Med (Mosk)*. 2003;81(11):35–38.
- Smith, K. A., Waypa, G. B., & Schumacker, P. T. (2017). Redox signaling during hypoxia in mammalian cells. *Redox Biology*, 13, 228–234.
- Smythies, J. A., Sun, M., Masson, N., Salama, R., Simpson, P. D., Murray, E., Neumann, V., Cockman, M. E., Choudhry, H., Ratcliffe, P. J., & Mole, D. R. (2019). Inherent DNA-binding specificities of the HIF-1α and HIF-2α transcription factors in chromatin. *EMBO Reports*, 20(1), e46401.
- Soliz, J., Joseph, V., Soulage, C., Becskei, C., Vogel, J., Pequignot, J. M., Ogunshola, O., & Gassmann, M. (2005). Erythropoietin regulates hypoxic ventilation in mice by interacting with brainstem and carotid bodies. *The Journal of Physiology*, 568(2), 559–571.
- Soliz, J., Thomsen, J. J., Soulage, C., Lundby, C., & Gassmann, M. (2009). Sex-dependent regulation of hypoxic ventilation in mice and humans is mediated by erythropoietin. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 296(6), R1837–R1846.
- Soo, J., Raman, A., Lawler, N. G., Goods, P. S. R., Deldicque, L., Girard, O., & Fairchild, T. J. (2023). The role of exercise and hypoxia on glucose transport and regulation. *European Journal of Applied Physiology*, 123(6), 1147–1165.
- Sorce, S., & Krause, K. H. (2009). NOX enzymes in the central nervous system: From signaling to disease. *Antioxidants & Redox Signaling*, 11(10), 2481–2504.
- Selfridge, A.C., Cavadas, M.A.S., Scholz, C.C., Campbell, E.L., Welch, L.C., Lecuona, E., Colgan, S.P., Barrett, K.E., Sporn, P.H.S., Szajner, J.I., et al. Hypercapnia suppresses the HIF-dependent adaptive response to hypoxia. *J. Biol. Chem.* 2016, 291, 11800–11808.
- Spaulding, H. R., & Yan, Z. (2022). AMPK and the adaptation to exercise. *Annual Review of Physiology*, 84(1), 209–227.
- Spliethoff, K., Meier, D., Aeberli, I., Gassmann, M., Langhans, W., Maggiorini, M., Lutz, T. A., & Goetze, O. (2013). Reduced insulin sensitivity as a marker for acute mountain sickness? *High Altitude Medicine & Biology*, 14(3), 240–250.
- Stewart, K.J. Role of exercise training on cardiovascular disease in persons who have type 2 diabetes and hypertension. *Cardiol. Clin.* 2004, 22, 569–586.
- Stray-Gundersen, S., Massoudian, S. D., Wojan, F., Tanaka, H., & Lalande, S. (2022). Hypoxic preconditioning reduces endothelial ischemia-reperfusion injury in older adults. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 323(5), R832–R838.
- Stuckless, T. J. R., Vermeulen, T. D., Brown, C. V., Boulet, L. M., Shafer, B. M., Wakeham, D. J., Steinback, C. D., Ayas, N. T., Floras, J. S., & Foster, G. E. (2020). Acute intermittent hypercapnic hypoxia and sympathetic neurovascular transduction in men. *The Journal of Physiology*, 598(3), 473–487.
- Suhovershin, A.V., Pantin, A.V., Suhovershin, R.A., Kisarova, Y.A., Kulikov, V.P., Bespalov, A.G. Restorative treatment of patients with neurasthenia with hypercapnic hypoxia in the conditions of a balneological resort. *Sibirskiy Vestnik Psichiatrii i Narkologii* =Siberian Bulletin of Psychiatry and Narcology. 2009;52 (13):126–128.

- Summers, B.A., Overholt, J.L., Prabhakar, N.R. CO<sub>2</sub> and pH independently modulate L-type Ca<sup>2+</sup> current in rabbit carotid body glomus cells. *J. Neurophysiol.* 2002, 88, 604–612.
- Sussman, M. A., Völkers, M., Fischer, K., Bailey, B., Cottage, C. T., Din, S., Gude, N., Avitabile, D., Alvarez, R., Sundararaman, B., Quijada, P., Mason, M., Konstandin, M. H., Malhowski, A., Cheng, Z., Khan, M., & McGregor, M. (2011). Myocardial AKT: The omnipresent nexus. *Physiological Reviews*, 91(3), 1023–1070.
- Syed, Z., Lin, H.-o-S., & Mateika, J. H. (2013). The impact of arousal state, sex, and sleep apnea on the magnitude of progressive augmentation and ventilatory long-term facilitation. *Journal of Applied Physiology*, 114(1), 52–65.
- Takeda, D., Hasegawa, T., Ueha, T., Imai, Y., Sakakibara, A., Minoda, M., Kawamoto, T., Minamikawa, T., Shibuya, Y., Akisue, T., et al. Transcutaneous carbon dioxide induces mitochondrial apoptosis and suppresses metastasis of oral squamous cell carcinoma in vivo. *PLoS ONE* 2014, 9, e100530.
- Takemori, T., Kawamoto, T., Ueha, T., Toda, M., Morishita, M., Kamata, E., Fukase, N., Hara, H., Fujiwara, S., Niikura, T., et al. Transcutaneous carbon dioxide application suppresses bone destruction caused by breast cancer metastasis. *Oncol. Rep.* 2018, 40, 2079–2087.
- Tan, A. Q., Barth, S., & Trumbower, R. D. (2020). Acute intermittent hypoxia as a potential adjuvant to improve walking following spinal cord injury: Evidence, challenges, and future directions. *Current Physical Medicine and Rehabilitation Reports*, 8(3), 188–198.
- Tao T., Liu Y., Zhang J. et al (2013) Therapeutic hypercapnia improves functional recovery and attenuates injury via antiapoptotic mechanisms in a rat focal cerebral ischemia/reperfusion model. *Brain Res* 1533:52–62.
- Taylor, C. T., & Colgan, S. P. (2017). Regulation of immunity and inflammation by hypoxia in immunological niches. *Nature Reviews Immunology*, 17(12), 774–785.
- Taylor, C. T., & Scholz, C. C. (2022). The effect of HIF on metabolism and immunity. *Nature Reviews Nephrology*, 18(9), 573–587.
- Tekin, D., Chou, E., Xi, L. Intermittent hypoxia and atherosclerosis. *Intermittent Hypoxia and Human Diseases*. 2012;29–45.
- Tekutyeva, N.A., Seroshtanova, E.V., Mikheeva, N.M., Vykhodtseva, G.I., Lobanov, Yu.F. The use of hypoxic-hypercapnic training on the respiratory simulator "Carbonic" in the treatment of neurogenic dysfunction of the bladder in children. *Fundamental'nye Issledovaniya =Fundamental Research*. 2014;10(2):371–374.
- Teppema, L. J. (2018). CrossTalk opposing view: The hypoxic ventilatory response does not include a central, excitatory hypoxia sensing component. *The Journal of physiology*, 596(15), 2939–2941.
- Teppema, L. J., & Dahan, A. (2010). The ventilatory response to hypoxia in mammals: Mechanisms, measurement, and analysis. *Physiological Reviews*, 90(2), 675–754.
- Terraneo, L., Paroni, R., Bianciardi, P., Giallongo, T., Carelli, S., Gorio, A., & Samaja, M. (2017). Brain adaptation to hypoxia and hyperoxia in mice. *Redox Biology*, 11, 12–20.
- Tester, N. J., Fuller, D. D., Fromm, J. S., Spiess, M. R., Behrman, A. L., & Mateika, J. H. (2014). Long-term facilitation of ventilation in humans with chronic spinal cord injury. *American Journal of Respiratory and Critical Care Medicine*, 189(1), 57–65.
- Timon, R., González-Custodio, A., Vasquez-Bonilla, A., Olcina, G., & Leal, A. (2022). Intermittent hypoxia as a therapeutic tool to improve health parameters in older adults. *International Journal of Environmental Research and Public Health*, 19(9), 5339.
- Tinkov, A.N., Aksenov, V.A. Effects of intermittent hypobaric hypoxia on blood lipid concentrations in male coronary heart disease patients. *High Alt Med Biol.* 2002;3(3):277–282.
- Tobin, B., Costalat, G., & Renshaw, G. M. (2020). Intermittent not continuous hypoxia provoked haematological adaptations in healthy seniors: Hypoxic pattern may hold the key. *European Journal of Applied Physiology*, 120(3), 707–718.
- Tobin, B., Costalat, G., & Renshaw, G. M. C. (2022). Pre-acclimation to altitude in young adults: Choosing a hypoxic pattern at sea level which provokes significant haematological adaptations. *European Journal of Applied Physiology*, 122(2), 395–407.
- Toriyama, T., Kumada, Y., Matsubara, T., Murata, A., Ogino, A., Hayashi, H., Nakashima, H., Takahashi, H., Matsuo, H., Kawahara, H. Effect of artificial carbon dioxide foot bathing on critical limb ischemia (fontaine IV) in peripheral arterial disease patients. *Int. Angiol.* 2002, 21, 367–373.
- Torres-Torrelo, H., Ortega-Sáenz, P., Gao, L., & López-Barneo, J. (2021). Lactate sensing mechanisms in arterial chemo-receptor cells. *Nature Communications*, 12(1), 4166.
- Tregub, P., Kulikov, V., Bespalov, A. Tolerance to acute hypoxia maximally increases in case of joint effect of normobaric hypoxia and permissive hypercapnia in rats. *Pathophysiology*. 2013;20(3):165–170.
- Tregub, P.P., Kulikov, V.P., Bespalov, A.G., Vvedensky, A.J., Osipov, I.S. Neuroprotective effects of individual or combined exposure to hypoxia and hypercapnia in the experiment. *Bull Exp Biol Med.* 2013;155(3):327–329.
- Tregub, P., Kulikov, V., Motin, Y. et al (2015) Combined exposure to hypercapnia and hypoxia provides its maximum neuroprotection effect during focal ischemic injury in the brain. *J Stroke Cerebrovasc Dis* 24(2):381–387.
- Tregub, P.P., Kulikov, V.P., Motin, Yu.G., Nagibaeva, M.E., Zabrodina, A.S. The stress of the endoplasmic reticulum of neurons in stroke is maximally limited by the combination of hypercapnia and hypoxia. *Bull Exp Biol Med.* 2016;161(4):457–461.
- Tregub, P.P., Kulikov, V.P., Stepanova, L.A., Zabrodina, A.S., Nagibaeva, M.E. The role of adenosine A1 receptors and mitochondrial K<sup>+</sup>-ATP channels in the mechanism of increasing resistance to acute hypoxia when combined with hypoxia and hypercapnia. *Patologicheskaya Fiziologiya i Eksperimental'naya Terapiya = Pathophysiology and Experimental Therapy*. 2014;58(4):48–52.

Tregub, P.P., Malinovskaya, N.A., Kulikov, V.P. et al (2016) Inhibition of apoptosis - a potential mechanism for the increased tolerance of brain ischemia with combination of hypercapnia and hypoxia. *Bull Exp Biol Med* 5:606–609. <https://doi.org/10.1007/s10517-016-3481-4>.

Tregub, P.P., Malinovskaya, N.A., Kulikov, V.P., Salmina, A.B., Nagibaeva, M.E., Zabrodina, A.S. et al. Inhibition of apoptosis is a potential mechanism of increasing ischemic tolerance of the brain with combined effects of hypercapnia and hypoxia. *Bull Exp Biol Med.* 2016;161(5):606–609.

Tregub, P.P., Malinovskaya, N.A., Osipova, E.D. et al (2022) Hypercapnia modulates the activity of adenosine A1 receptors and mitoK<sub>ATP</sub>-channels in rat brain when exposed to intermittent hypoxia. *Neuromol Med* 24(2):155–168. <https://doi.org/10.1007/s12017-021-08672-0>.

Trumbower, R. D., Jayaraman, A., Mitchell, G. S., & Rymer, W. Z. (2012). Exposure to acute intermittent hypoxia augments somatic motor function in humans with incomplete spinal cord injury. *Neurorehabilitation and Neural Repair*, 26(2), 163–172.

Tsarev, A.Yu., Kunitsyna, L.A., Yezhov, V.V., Kolesnikova, E.Yu., Platunova, T.E., Chernysh, D.A. The dynamics of external respiration, vegetative and cognitive functions under the influence of normobaric hypoxic-hypercapnic training in patients with chronic cerebral ischemia of atherosclerotic genesis. *Vestnik Phizioterapii i Kurortologii = Bulletin of Physiotherapy and Balneology*. 2017;23(1):28–33.

Ueha, T., Oe, K., Miwa, M., Hasegawa, T., Koh, A., Nishimoto, H., Lee, S., Niikura, T., Kurosaka, M., Kuroda, R., et al. Increase in carbon dioxide accelerates the performance of endurance exercise in rats. *J. Physiol. Sci.* 2018, 68, 463–470.

Ure, B.M., Niewold, T.A., Bax, N.M.A., Ham, M., van de Zee, D.C., Essen, G.J. Peritoneal, systemic, and distant organ inflammatory responses are reduced by a laparoscopic approach and carbon dioxide vs air. *Surg. Endosc.* 2002, 16, 836–842.

Vadász, I., Dada, L.A., Briva, A., Trejo, H.E., Welch, L.C., Chen, J., Tóth, P.T., Lecuona, E., Witters, L.A., Schumacker, P.T., et al. AMP-activated protein kinase regulates CO<sub>2</sub>-induced alveolar epithelial dysfunction in rats and human cells by promoting Na<sub>K</sub>-ATPase endocytosis. *J. Clin. Investig.* 2008, 118, 752–762.

Van Meijel, R. L. J., Vliex, L. M. M., Hartwig, S., Lehr, S., Al-Hasani, H., Blaak, E. E., & Goossens, G. H. (2023). The impact of mild hypoxia exposure on myokine secretion in human obesity. *International Journal of Obesity*, 47(6), 520–527.

Vannucci, R.C., Towfighi, J., Heitjan, D.F., Brucklacher, R.M. Carbon dioxide protects the perinatal brain from hypoxic-ischemic damage: an experimental study in the immature rat. *Pediatrics*. 1995;95(6):868–874.

Veglio, M., Maule, S., Cametti, G., Cogo, A., Lussiana, L., Madrigale, G., & Pecchio, O. (1999). The effects of exposure to moderate altitude on cardiovascular autonomic function in normal subjects. *Clinical Autonomic Research: Official Journal of the Clinical Autonomic Research Society*, 9(3), 123–127.

Velizhanina, I.A., Evdokimova, O.A., Gapon, .L.I., Vdovenko, S.V. The response to normoxic hypoxia and its duration in arterial hypertension patients as shown by a prospective study. *Klin Med (Mosk)*. 2002;80(5):22–26.

Vermeulen, T. D., Benbaruj, J., Brown, C. V., Shafer, B. M., Floras, J. S., & Foster, G. E. (2020). Peripheral chemoreflex contribution to ventilatory long-term facilitation induced by acute intermittent hypercapnic hypoxia in males and females. *The Journal of Physiology*, 598(20), 4713–4730.

Veselá, A., Wilhelm, J. The role of carbon dioxide in free radical reactions of the organism. *Physiol. Res.* 2002, 51, 335–339.

Vinitskaia, R.S., Davidov, E.G., Struchkov, P.V. Hypoxic and hypercapnic gas mixtures in complex treatment and rehabilitation of patients with chronic obstructive diseases. In: *Intermittent Hypoxic Training: Effectiveness and Mechanisms of Action*. Institute of Physical Culture, Kiev, 1992. P. 62–64.

Volkova, Y. L., Pickel, C., Jucht, A. E., Wenger, R. H., & Scholz, C. C. (2022). The asparagine hydroxylase FIH: A unique oxygen sensor. *Antioxidants & Redox Signaling*, 37(13–15), 913–935.

Wadhwa, H., Gradinaru, C., Gates, G. J., Badr, M. S., & Mateika, J. H. (2008). Impact of intermittent hypoxia on long-term facilitation of minute ventilation and heart rate variability in men and women: do sex differences exist? *Journal of Applied Physiology*, 104(6), 1625–1633.

Wakhloo, D., Scharkowski, F., Curto, Y., Javed Butt, U., Bansal, V., Steixner-Kumar, A. A., Wüstefeld, L., Rajput, A., Arinrad, S., Zillmann, M. R., Seelbach, A., Hassouna, I., Schneider, K., Qadir Ibrahim, A., Werner, H. B., Martens, H., Miskowiak, K., Wojcik, S. M., Bonn, S., Nacher, J., Nave, K.-A., & Ehrenreich, H. (2020). Functional hypoxia drives neuroplasticity and neurogenesis via brain erythropoietin. *Nature Communications*, 11(1), 1313.

Walter, P., & Ron, D. (2011). The unfolded protein response: From stress pathway to homeostatic regulation. *Science* (New York, NY), 334(6059), 1081–1086.

Wang, G. L., Jiang, B. H., Rue, E. A., & Semenza, G. L. (1995). Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O<sub>2</sub> tension. *Proceedings of the National Academy of Sciences, USA*, 92(12), 5510–5514.

Wang, H., Shi, X., Schenck, H., Hall, J. R., Ross, S. E., Kline, G. P., Chen, S., Mallet, R. T., & Chen, P. (2020). Intermittent hypoxia training for treating mild cognitive impairment: A pilot study. *American Journal of Alzheimer's Disease and Other Dementias*, 35(17), 153331751989672.

Wang, Z., Su, F., Bruhn, A., Yang, X., Vincent, J. Acute Hypercapnia Improves Indices of Tissue Oxygenation More than Dobutamine in Septic Shock. *Amer J Resp and Crit Care Med* 2008;177:178-183.

West J.B. High altitude medicine. *Am J Respir Crit Care Med*. 2012;186(12):1229–1237.

Westermaier, T., Stetter, C., Kunze, E., Willner, N., Holzmeier, J., Kilgenstein, C., et al. Controlled transient hypercapnia: a novel approach for the treatment of delayed cerebral ischemia after subarachnoid hemorrhage? *J Neurosurg*. 2014;121(5):1056–1062.

Willie, C. K., Tzeng, Y.u-C., Fisher, J. A., & Ainslie, P. N. (2014). Integrative regulation of human brain blood flow. *The Journal of Physiology*, 592(5), 841–859.

- Wollina, U., Heinig, B., Uhlemann, C. Transdermal CO<sub>2</sub> application in chronic wounds. *Int. J. Low. Extrem. Wounds* 2004, 3, 103–106.
- Wu, J., Li, P., Wu, X., & Chen, W. (2015). Chronic intermittent hypoxia decreases pain sensitivity and increases the expression of HIF1α and opioid receptors in experimental rats. *Sleep & Breathing = Schlaf & Atmung*, 19, 561–568.
- Xing, T., Pilowsky, P. M., & Fong, A. Y. (2014). Mechanism of sympathetic activation and blood pressure elevation in humans and animals following acute intermittent hypoxia. *Progress in Brain Research*, 209, 131–146.
- Xu, Y.-J., Elimban, V., Bhullar, S.K., Dhalla, N.S. Effects of CO<sub>2</sub> water-bath treatment on blood flow and angiogenesis in ischemic hind limb of diabetic rat. *Can. J. Physiol. Pharmacol.* 2018, 96, 1017–1021.
- Xu, Y., Elimban, V., Dhalla, N.S. Carbon dioxide water-bath treatment augments peripheral blood flow through the development of angiogenesis. *Can. J. Physiol. Pharmacol.* 2017, 95, 938–944.
- Yamada, M., Ishikawa, T., Yamanaka, A., Fujimori, A., Goto, K. Local neurogenic regulation of rat hindlimb circulation: CO<sub>2</sub>- induced release of calcitonin gene-related peptide from sensory nerves. *Br. J. Pharmacol.* 1997, 122, 710–714.
- Yan, W., Chen, Z., Chen, J., & Chen, H. (2016). Isoflurane preconditioning protects rat brain from ischemia reperfusion injury via up-regulating the HIF-1alpha expression through Akt/mTOR/s6K activation. *Cellular and Molecular Biology*, 62, 38–44.
- Yan, Z., Okutsu, M., Akhtar, Y.N., Lira, V.A. Regulation of exercise-induced fiber type transformation, mitochondrial biogenesis, and angiogenesis in skeletal muscle. *J. Appl. Physiol.* 2011, 110, 264–274.
- Yang C.C., Lin L.C., Wu M.S. et al (2009) Repetitive hypoxic preconditioning attenuates renal ischemia/reperfusion induced oxidative injury via upregulating HIF-1 alpha-dependent Bcl-2 signaling. *Transplantation* 88:1251–1260.
- Yatagai, N., Hasegawa, T., Amano, R., Saito, I., Arimoto, S., Takeda, D., Kakei, Y., Akashi, M. Transcutaneous carbon dioxide decreases immunosuppressive factors in squamous cell carcinoma. *In Vivo. BioMed Res. Int.* 2021, 2021, 1–9.
- Yuan, G., Khan, S. A., Luo, W., Nanduri, J., Semenza, G. L., & Prabhakar, N. R. (2011). Hypoxia-inducible factor 1 mediates increased expression of NADPH oxidase-2 in response to intermittent hypoxia. *Journal of Cellular Physiology*, 226(11), 2925–2933.
- Zakynthinos S., Katsaounou P., Karatza M.H. et al (2007) Antioxidants increase the ventilatory response to hyperoxic hypercapnia. *Am J Respir Crit Care Med* 175:62–68.
- Zenewicz, L. A. (2017). Oxygen Levels and Immunological Studies. *Frontiers in Immunology*, 8, 324.
- Zhang, J., Haddad, G. G., & Xia, Y. (2000). delta-, but not mu- and kappa-, opioid receptor activation protects neocortical neurons from glutamate-induced excitotoxic injury. *Brain research*, 885(2), 143–153.
- Zhang, J.-M., & An, J. (2007). Cytokines, inflammation, and pain. *International Anesthesiology Clinics*, 45(2), 27–37.
- Zhao, H., Ren, C., Chen, X., & Shen, J. (2012). From rapid to delayed and remote postconditioning: The evolving concept of ischemic postconditioning in brain ischemia. *Current drug targets*, 13(2), 173–187.
- Zhao Z.S., Khan S., O'Brien P.J. (1998) Catecholic iron complexes as cytoprotective superoxide scavengers against hypoxia: reoxygenation injury in isolated hepatocytes. *Biochem Pharmacol* 56(7):825–830. [https://doi.org/10.1016/s0006-2952\(98\)00222-6](https://doi.org/10.1016/s0006-2952(98)00222-6).
- Zhou Q., Cao B., Niu L. et al (2010) Effects of permissive hypercapnia on transient global cerebral ischemia-reperfusion injury in rats. *Anesthesiology* 112:288–297